

Comparative Study on Malondialdehyde and Certain Antioxidants in North West obese Indians

Kuldip Singh^{1*} and Saranpal Singh²

¹Department of Biochemistry, Govt. Medical College Amritsar (Punjab), India, Presently working in Department of Biochemistry, Govt. Medical College Patiala (Punjab), India.

²Department of Biochemistry, Adesh Institute of Medical Sciences-Bathinda (Punjab), India.

ABSTRACT

Background: Obesity is one of the most common health problems in developed and developing nations like India. In addition, obesity is considered as a principal risk factor in the initiation of various non communicable chronic diseases such as dyslipidemia, atherosclerosis, cardiovascular, non-insulin dependent diabetes mellitus etc. The pathogenesis of these diseases is associated with oxygen-derived free radicals. **Aim:** The aim of the study was to analyze oxidative status in obese subjects with respect to normal healthy subjects. **Material and Methods:** The present study was performed in 50 obese men (BMI>46 kg/m²) and equal number of normal healthy men (BMI< 27 kg/m²) and the levels of malondialdehyde, antioxidant enzymes/molecules like superoxide dismutase, catalase reduced glutathione, glutathione reductase, glutathione peroxidase, antioxidant Vitamin-E and Vitamin-C along with total antioxidant activity were evaluated. **Results:** A significant increase (p<0.01) in the concentration of malondialdehyde by 61.07% was observed in obese men subjects in comparison to normal healthy men subjects while a significant fall was recorded in the levels of vitamin-E (19.54%, p<0.05), Vitamin-C (18.07%, p<0.05), superoxide dismutase (19.51%, p<0.05), catalase (20.28%, p<0.05), GSH (36.56%, p<0.05), GR (33.28%, p<0.01), GPx (19.14%, p<0.05) and a significant decrease by 33.33% (p<0.01) was observed in the total antioxidant activity in obese subjects as compared to normal healthy control subjects. **Conclusion:** Aforementioned observations suggested that oxidative is induced in the obese subjects (BMI>46 kg/m²) due to alterations in the malondialdehyde and other oxidative stress markers such as Vitamin-E, Vitamin-C, Superoxide dismutase, catalase, reduced glutathione, glutathione reductase and glutathione peroxidase might be responsible for the etiology of diseases like diabetes, cancer arthritis, atherosclerosis etc. and therefore, obesity should be treated with the same attention of non-communicable diseases like atherosclerosis, arthritis, diabetes, cancer etc.

Key words: Obesity, Oxidative stress, Vitamin-E, Vitamin-C, Lipoproteins, Lipid peroxidation, Antioxidants, Cardiovascular disease (CVD).

INTRODUCTION

Obesity is a growing epidemic in developed and developing nations like India, with the estimated prevalence ranging from 10 to 50 percent or more in the adult population.¹ Many studies had confirmed the role of obesity in diabetes mellitus, hyperlipidemia, colon cancer, sudden death and other cardiovascular diseases has been confirmed.²⁻⁴ It is

also well reported in the literature that oxidative stress is responsible for the pathophysiology of all these disease.^{3,5-7} Many cellular processes such as control of membrane permeability, secretion of extra cellular matrix proteins, cell proliferation, cell migration and cell deaths are found to be promoted by oxygen free radicals that have contributed to the development of diabetes and cancer.⁵ Free radicals may produce their deleterious effects by; a) Deoxidizing cell membrane lipids thus altering the structural integrity of the membrane; b) Peroxidizing the lipids in maintaining the micro environment of membrane proteins or reacting both with proteins and with membrane lipids separately or concurrently.

*Corresponding address:

Dr. Kuldip Singh

Associate Professor, Department of Biochemistry,
Govt. Medical College-Patiala (Punjab), India.

E-mail: drkuldip08@gmail.com

DOI: 10.5530/jcdr.2015.3.5

In addition, oxygen radicals and products of the reaction may alter the structure of DNA and RNA. It is apparent therefore that an excessive stress imposed upon the cell could lead to the cellular modifications and thereby, essential chemical components of the cells leading to alter function. The cell and plasma contain a variety of antioxidants and free radical scavenging molecules and enzymes like Superoxide dismutase (SOD), Catalase (CAT), Xanthine oxidase (XOD), Glutathione (GSH), Glutathione reductase (GR), Glutathione peroxidase (GPx) and Glutathione-S-Transferase (GST) etc., which under normal circumstances help the cell to maintain a reducing environment by preventing the potentially deleterious effects of free radical on cell membrane and organelles.⁸ Any changes in the activity of these enzymes and antioxidants increase oxidative stress and could lead to the development of various diseases like cardiovascular diseases, arthritis, diabetes, cancer etc.

Although the exact biochemical mechanisms responsible for the association between obesity and the development of various non-communicable diseases like atherosclerosis, arthritis, diabetes, cancer etc. have not been completely elucidated, and data on the change in antioxidant molecules/enzymes such as SOD, CAT, GR, GPx, GSH is scant in obese North punjabi population of Indian. So, the present work was designed, to study the change in oxidative stress markers by analyzing MDA, SOD, CAT, GSH GR, GPx along with change in antioxidant vitamins such as Vitamin-E and Vitamin-C levels in obese North Punjabi population of India.

MATERIAL AND METHODS

The present study was carried out in the Department of Biochemistry, Adesh Medical College Bathinda on 50 male obese individuals in age group ranging from 20 to 35 years with BMI > 46 kg/m². The obese subjects, who were suffered with complication such as diabetes mellitus, hypertension, hypothyroidism, renal failure, hepatic diseases, acute illnesses, recurrent myocardial infarction, unstable angina and those not on any weight lose treatment and inherited obese individuals were not taken for the study and equal number of normal healthy male subjects in same age group ranging from 20 to 35 years with BMI < 27 kg/m² were taken as control group.

All subjects recruited for the study were vegetarian, non-smoker and non-alcoholic, and there was no positive family history of diabetes, Cardiovascular diseases (CVD) in these subjects. These subjects were interviewed and by giving a questionnaire regarding the detailed information on their

lifestyle, medical history, diet etc. and after obtaining the written consent these were considered for the study.

Study design

This study was a case control cross-sectional prospective study. This study protocol was approved by the Ethical Committee of the Institution. A detailed history, physical, and systemic examination, including measurement of height, weight, heart rate, blood pressure, and Body mass index (BMI), complete lipid profile, fasting and post-prandial glucose levels was carried out in every subject who entered the study as per a pre-designed proforma for assessing the signs of chronic heart failure, diabetes and also the presence of any exclusion criteria.

DATA COLLECTION

Measurements of Anthropometric Parameters

The examination of body weight was done by taking weight in kilogram (kg) and height was measured in centimeters. The BMI was calculated from the formula: BMI = weight in kg / (height in meters)².

Blood sampling

A volume of 5 ml of peripheral venous blood was collected by vein puncture using a dry, disposable syringe between 8 and 9 AM after an overnight fast from both groups (control and Obese patients) in heparin containing vial. Blood samples were immediately centrifuged at 4000 rpm at 4°C and plasma was stored at -20°C until analysis. The 100 µl of RBC were taken and lysed with 2.0 ml ice cold water and the clear lysate obtained after spinning down the cell debris at 8500 g for 10 min at 4°C was used for various biochemical assays.

Biochemical Assays

Estimation of Glucose levels

Fasting and post prandial glucose levels were estimated spectrophotometrically using an enzymatic test kit based on GOD-POD method supplied by Transasia Biomedical Private Limited, Mumbai (India).

Estimation of Lipoprotein levels

The total cholesterol, triglyceride, high density lipoprotein cholesterol, were analyzed by using commercial available kits procured from Transasia Biomedical Private Limited, Mumbai (India). Low density lipoprotein-cholesterol levels were estimated by the primary measurements by using the empirical equation of Bates and Warren, 1989.⁹ Serum LDL-cholesterol = Total cholesterol - [HDL + Triglyceride/5] and

Table 1: General characteristics of normal healthy control and obese population

Anthropometric and Biochemical Assays	Normal Healthy Control Subjects	Obese Subjects
Subject Number	50	50
Sex	Male	Male
Height (cm)	158.12 ± 0.16	161.6 ± 0.1
Weight (kg)	67.00 ± 14	121.00 ± 25
Age (years)	31.17 ± 5.45	33.34 ± 6.51
Body mass index (Kg/m ²)	26.80 ± 0.48	46.33 ± 0.40
Blood pressure systolic (mmHg)	128.15 ± 10.53	127.45 ± 12.53
Blood pressure diastolic (mmHg)	85.01 ± 7.32	85.29 ± 8.74
Hemoglobin (g/dl)	14.81 ± 1.16	13.89 ± 2.91
Fasting Glucose (mg/dl)	84.56 ± 9.23	89.65 ± 11.21
Postprandial Glucose(mg/dl)	121.16 ± 9.11	124.00 ± 8.29
Total Cholesterol (mg/dl)	140.88 ± 28.92	158.70 ± 34.57
Triglycerides (mg/dl)	80.01 ± 18.65	86.06 ± 24.62
LDL-Cholesterol (mg/dl)	94.85 ± 15.40	96.09 ± 16.05
HDL-Cholesterol (mg/dl)	45.80 ± 14.60	48.09 ± 18.70
VLDL-Cholesterol (mg/dl)	16.00 ± 11.41	17.01 ± 14.65

very low density lipoprotein cholesterol was calculated by dividing triglyceride concentration with 5.

Estimation of hemoglobin (%)

Hemoglobin (Hb) from whole blood was estimated spectrophotometrically by using a kit supplied by Transasia Biomedical Private Limited, Mumbai (India).

Estimation of malondialdehyde (MDA) levels

MDA level, an end product of lipid peroxidation of erythrocytes was assayed using a diagnostic kit supplied by Transasia Biomedical Private Limited, Mumbai (India).

- **Estimation of Superoxide dismutase (EC 1.15.1.1):** SOD levels were measured in erythrocyte.¹⁰
- **Estimation of Catalase (EC 1.11.1.6):** CAT activity was measured spectrophotometrically¹¹ in erythrocyte by following the decomposition rate of H₂O₂ at 240 nm.
- **Estimation of Reduced Glutathione (GSH):** The levels of GSH was estimated in whole blood.¹²
- **Estimation of Glutathione peroxidase (EC 1.11.1.9):** The activity of GPx in erythrocytes was estimated spectrophotometrically.¹³
- **Estimation of Glutathione reductase (1.6.4.2):** The GR activity was determined in Erythrocyte.¹⁴
- **Estimation of Vitamin C:** Vitamin C was estimated in Plasma.¹⁵ Vitamin C is oxidized to diketogluconic acid which reacts with 2, 4-dinitrophenyl hydrazine to form diphenylhydrazone. The hydrazone dissolves in strong acid solution to form orange-red colored complex at 520 nm. Vitamin C was expressed in mg/dl.
- **Estimation of Vitamin E:** Vitamin E was estimated in Plasma.¹⁶ Vitamin E reduces ferric ion to ferrous

ion, which forms a red colored complex with α , α '-bipyridyl. Vitamin E was expressed in mg/dl.

- **Estimation of Total Antioxidant Activity (TAA):** Plasma total antioxidant capacity was estimated in plasma by the FRAP (Ferric Reducing Ability of Plasma) assay.¹⁷

Statistical Analysis

The data was expressed as Mean ± SD and analyzed with the SPSS 16.0.7 statistical software package. Differences between the obese and control subjects were evaluated using the Student's independent samples *t* test. Differences were considered statistically significant at *P*<0.05.

RESULTS

- **Anthropometric Parameters:** The Anthropometric measurements of both obese and normal healthy control subjects are summarized in the Table 1. The body weight, height, BMI, BP systolic and BP-diasystolic was 137 ± 25 Kg, 236.6 ± 0.11 cm, 46.33 ± 0.40 Kg/m², 127.45 ± 12.53 mmHg, 85.29 ± 8.74 mmHg respectively in obese subjects in comparison to 62 ± 14 Kg, 249.12 ± 0.16 cm, 26.80 ± 0.48 Kg/m², 128.15 ± 10.53 mmHg and 85.01 ± 7.32 mmHg respectively of healthy control subjects.
- **Glucose, Hemoglobin and Lipid levels:** The levels of fasting and post prandial glucose levels, hemoglobin and lipoprotein fractions are summarized in Table 1. The fasting and post prandial plasma glucose levels was 89.65 ± 11.21 mg/dl and 124 ± 8.29 mg/dl respectively in obese patients with respect to 84.56 ± 9.23 mg/dl and 121.16 ± 9.11 mg/dl fasting and post prandial glucose levels in healthy control subjects. A hemoglobin level 13.89 ± 2.91 g/dl in obese subjects and 14.81 ±




Figure 1: Changes in MDA, SOD, CAT, GSH, GR and GPx in normal healthy controls subjects (n=50) and obese patients (n=50)

Values in figure are expressed as mean ± standard deviation, The levels of MDA are expressed in nmol/g Hb, SOD levels are expressed in U/g Hb, and levels of CAT, GSH, GR & GPx are expressed in U/g Hb, mg/g Hb, mU/g Hb, and mU/g Hb respectively.




Figure 2: Alterations in vitamin-E and Vitamin-C levels in normal healthy controls subjects (n=50) and obese patients (n=50)

Values in figure are expressed as mean ± standard deviation; The levels of vitamin -E and Vitamin-C are expressed in mg/dl.

1.16 g/dl in normal healthy subjects was observed. The lipoprotein fractions such as total cholesterol, triglycerides, LDL-cholesterol, HDL-cholesterol and VLDL-cholesterol levels was 178.70 ± 34.57 mg/dl, 156.96 ± 24.62 mg/dl, 109.5 ± 20.6 mg/dl, 38 ±

18.70 mg/dl, 31.8 ± 14.65 mg/dl in obese patients in comparison to 140.88 ± 28.92 mg/dl, 104.45 ± 18.65 mg/dl, 94.85 ± 15.40 mg/dl, 45.8 ± 14.60 mg/dl and 21.1 ± 11.41mg/dl total cholesterol, triglycerides, LDL-cholesterol, HDL- cholesterol and VLDL- cholesterol




Figure 3: Changes in total antioxidant activity in normal healthy controls subjects (n = 50) and obese patients (n = 50).
Values in figure are expressed as mean ± standard deviation. The levels of total antioxidant activity are expressed in µ mol/L.

respectively in normal healthy control subjects. There were statistical significant difference in the glucose, hemoglobin and various lipid levels in obese and normal healthy subjects and these biochemical parameters were remained within the normal limit.

- **Lipid Peroxidation:** A significant increase was recorded in the MDA levels (from 132.93 ± 11.23 to 214.12 ± 12.27 nmol MDA/gHb) in obese subjects by 61.07% ($p < 0.001$) with respect to normal healthy subjects (Figure 1).
- **Antioxidants vitamin and Enzymes:** A significant decrease ($p < 0.05$) was seen in plasma Vitamin-E (From 0.87 ± 0.16 to 0.70 ± 0.10 mg/dl) and Vitamin-C (From 0.83 ± 0.06 to 0.68 ± 0.06 mg/dl) by 19.54% and 18.07% respectively in obese with respect to normal healthy subjects (Figure 2). A significant decrease in the levels of SOD (from 239.52 ± 8.37 to 192.78 ± 6.72 U/g Hb), CAT (from 81.4 ± 2.71 to 69.89 ± 4.19 mg/g Hb), GSH (71.55 ± 1.92 to 35.39 ± 2.14 mg/g Hb), GR (from 45.4 ± 3.16 to 30.29 ± 3.62) and GPx (from 39.49 ± 2.61 to 32.93 ± 3.92 mg/gHb) by 19.51% ($p < 0.05$), 20.28% ($p < 0.05$), 36.56% ($p < 0.001$), 33.28% ($p < 0.001$), 19.14% ($p < 0.05$) respectively was observed in erythrocyte of obese patients in comparison to normal healthy control subjects (Figure 1).
- **Total Antioxidant Activity:** The levels of Total Antioxidant Activity was found 799.26 ± 19.69 µ mol/L in plasma of obese individuals and $613.23 \pm$

11.07 µ mol/L in normal healthy control individuals. This fall in plasma total antioxidant activity by 23.30% was found to be statistically significant ($p < 0.01$) in obese subjects with respect to control healthy subjects (Figure 3).

DISCUSSION

In the present case control study, we recorded a significant increase ($p < 0.001$) by 61.07% in MDA (representing the lipid peroxidation) levels in obese individuals even in the absence of any chronic diseases such as diabetes, hypertension, hyperlipidemia etc in comparison to normal healthy individuals (Figure 1). Lipid peroxidation causes polymerization of membrane components and induce changes in the membrane permeability resulting in hemolysis, would relate to the degree of intravascular Red blood cell (RBC) destruction. Extravascular mechanisms of RBC destruction may involve changes in cell deformability and antigenicity. Our observation of increased malondialdehyde levels is agreement with the literature reports and to decrease the membrane fluidity, deformability, visco-elasticity and life span of erythrocytes in the obese subjects,¹⁸⁻²¹ which might be responsible for the initiation of complications such as diabetes, hypertension, hyperlipidemia etc. in obese subjects later on in the north western Indian population.

The biological effects of free radicals are normally controlled *in vivo* by a wide range of antioxidants such as vitamin E, the main liposoluble chain breaking antioxidant

in human beings, scavenges peroxy radicals produced during lipid peroxidation. Reduced glutathione and vitamin C regenerate vitamin E. A significant decrease in vitamin-E and vitamin-C, have potential to protect both cytosolic and membrane components of cells from oxidant damage²¹ levels were falls significantly in the plasma of obese subjects with respect to normal healthy subjects (Figure 2) indicated the accumulation of free radicals.

SOD, a superoxide radical scavenging enzyme is considered the first line of defense against the deleterious effect of oxygen radicals in the cells and it scavenges reactive oxygen radical species by catalyzing the dismutation of O_2^- radical to H_2O_2 and O_2 .⁵⁻⁸ A significant inhibition by 25.63% in SOD activity in diabetics (Figure 1) may results in an increased flux of O_2^- radical and hence reflects the tissue damage/injury. The activity of CAT, another potent antioxidant enzyme, especially against the O_2^- radicals and singlet oxygen, was also found to significantly ($p < 0.01$) decreased by 20.28% in erythrocyte of obese subjects by with respect to control subjects (Figure 1) CAT, protects cells from the accumulation of H_2O_2 by dismutating it to form H_2O and O_2 or by using it as an oxidant, in which it works as a peroxidase.^{6,22} Therefore, the decrease in the activity of CAT suggested the accumulation of H_2O_2 in the obese subjects.

Glutathione (GSH), a tripeptide is maintained in reduced state by an efficient glutathione peroxidase/glutathione reductase system. Glutathione is a potent endogenous antioxidant that helps to protect cells from a number of noxious stimuli including oxygen derived free radicals.^{7,23,24} In the present work, the level of glutathione significantly decreased by 36.56% ($p < 0.01$) in obese subjects with respect to normal healthy control subjects (Figure 1). A significant fall in GSH levels, confirm an increased susceptibility to oxidative damage. The activity of GPx, a selenium-containing enzyme was found to be decreased by 19.14% ($p < 0.05$) in obese individuals in comparison to control healthy individuals (Figure 1). GPx catalyses the reduction of variety of hydrogen peroxide ($ROOH$ and H_2O_2) using glutathione as a substrate, thereby protecting mammalian cells against oxidative stress.²⁵ The significant inhibition ($p < 0.001$, 44.53%) in the activity of GR in diabetics (Figure 1) attributed to increased oxidation or decreased synthesis of GSH. The less availability of NADPH may also cause a decrease in GR activity.²⁶ Our observations of significant reduction in GSH levels are in an agreement with the literature reports that inverse relationship exists between LPO and GSH status, can impair the cell defense against the toxic action of xenobiotic and may lead to cell injury/

death.²⁴ A significant fall in GPx activity may render the tissue more susceptible to LPO damage. Accordingly, in the present work, we observed a significant decrease in GPx activity upon increase in LPO level. This observation is in accordance with the hypothesis that LPO and GPx might play a role in tissue damage²⁷⁻²⁹ that oxidative stress is induced in the obese subjects (Figure 3).

CONCLUSION

Aforementioned observations suggested that oxidative is induced in the obese subjects due to alterations in the levels of MDA, vit-E, vit-C, SOD, CAT, GSH, GR, GPx might be responsible for the pathophysiology of diseases like diabetes, cancer arthritis, atherosclerosis etc. and therefore, obesity should be treated with the same attention of various non-communicable diseases like atherosclerosis, arthritis, diabetes, cancer etc. Further studies using antioxidant enzymes, vitamin supplementations alone and in combinations, lifestyle and dietary modification might be beneficial to reduce oxidative stress in obese individuals.

ACKNOWLEDGEMENT

Authors are extremely thankful to the participants for the completion of this work.

ABBREVIATIONS

CVDs	: Cardiovascular disease
DM	: Diabetes Mellitus
MDA	: Malondialdehyde
LPO	: lipid peroxidation
SOD	: Superoxide dismutase
CAT	: Catalase
GSH	: Reduced Glutathione
GR	: Glutathione Reductase
GPx	: Glutathione Peroxidase
TAA	: Total Antioxidant Activity
Vit.-E	: Vitamin-E
Vit.-C	: Vitamin-C
H_2O	: Water
O_2	: Oxygen
H_2O_2	: Hydrogen Peroxide
O_2^-	: Superoxide anion.

REFERENCES

- Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes.* 2008; 32(9): 1431-7.
- Olusi SO. Obesity is an independent risk factor for plasma lipid peroxidation and depletion of erythrocyte cytoprotective enzymes in humans. *Int J Obes.* 2002; 26(9): 1159-64.
- Skalicky J, Muzakova V, Kandar R, Meloun M, Rousar T. Oxidative stress and metabolic syndrome in obese adults with and without controlled diet restrictions. *Bratisl Lek Listy.* 2009; 110(3): 152-7.
- Selvakumar C, Uma MS, Archana L. Oxidant-Antioxidant disturbance in men classified as obese according to the preliminary WHO guidelines for Asians. *J Stress Physiol and Biochem.* 2012; 8(1): 172-81.
- Kuldip Singh. Role of reactive oxygen species in health and diseases. Edited by Lap Lambert Academic Publishing Germany; 2014. p. 1-60.
- Kuldip Singh, Ahluwalia P. Effect of oral ingestion of monosodium glutamate administration on various lipid fractions and certain antioxidant enzymes in cardiac tissue of chronic alcoholic adult male rats. *J Cardiovas Des. Res.* 2012; 3(1): 215-21.
- Kuldip Singh, Saranpal Singh, Seema Chopra. Ameliorative effect of lycopene on lipid peroxidation and certain antioxidant activity in diabetic patients. *J Diabetes and Metab.* 2012; 3(6): 2-5.
- Sukhjeet K, Kuldip S. Comparative study on MDA and SOD in IHD patients and healthy smokers edited by Lap Lambert Academic Publishing Germany; 2014. p. 1-210.
- Bates CM, Warren GS. Estimation of low density lipoprotein cholesterol. *South Med. J.* 1989; 82(1): 570-5.
- Jaiswal D, Rai PK, Mehta S, Chatterji S, Shukla S, Rai DK, *et al.* Role of *Moringa oleifera* in regulation of diabetes-induced oxidative stress. *Asian Pac J Trop Med.* 2013; 6(6): 426-32.
- Al-Essa M, Dhaunsi GS, Al-Qabandi W, Khan I. Impaired NADPH oxidase activity in peripheral blood lymphocytes of galactosemia patients. *Exp Biol Med.* 2013; 238(7): 779-86.
- Giustarini D, Dalle-Donne I, Milzani A, Fanti P, Rossi R. Analysis of GSH and GSSG after derivatization with N-ethylmaleimide. *Nat Protoc.* 2013; 8(9): 1660-9.
- Worthington DJ, Rosemeyer MA. Glutathione reductase from human erythrocytes. Molecular weight, subunit composition and aggregation properties. *Eur J Biochem.* 1975; 60(4): 459-66.
- Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med.* 1967; 70(1): 158-69.
- Natelson S. In *Techniques in Clinical Biochemistry*. CC Thomas, USA 1971; 162: 288.
- Baker H, Frank O. *Clinical Vitaminology, methods and interpretations.* Inter science Publishers. John Wiley and Sons Inc., New York; 1968. 172.
- Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power-the FRAP assay. *Anal Biochem.* 1996; 239(1): 70-6.
- Cazzola R, Rondanelli M, Russo-Volpe S, Ferrari E, Castaro B. Decreased membrane fluidity and altered susceptibility to peroxidation and lipid composition in overweight and obese female erythrocytes. *J Lipid Res.* 2004; 45(10): 1846-51.
- Kelishadi R, Hashemipour M, Sheikh HA, Samani SG. Changes in serum lipid profile of obese or overweight children and adolescents following a lifestyle modification course. *ARYA Atheroscler.* 2012; 8(3): 143-8.
- Amirkhizi F, Siassi F, Minaie S, Djalali M, Rahimi A, Chamari M. Is obesity associated with increased plasma lipid peroxidation and oxidative stress in women?. *ARYA Atherosclerosis Journal* 2007; 2(4): 189-92.
- Madhikarmi NL, Murthy KRS, Rajagopal V, Singh PP. Lipid peroxidation and antioxidant status in patients with type 2 diabetes in relation to obesity in Pokhara-Nepal. *J Diabetol.* 2013; 1(3): 3-8.
- Kuldip Singh, Jaspinder S, Ahluwalia P. Effect of oral ingestion of monosodium glutamate administration on various lipid fractions, glutathione and its metabolizing enzymes in arterial tissue of chronic alcoholic adult male rats. *Toxicol. International* 2012; 18(2): 33-8.
- Kuldip Singh. Alterations in serum glutathione as a marker of oxidative stress in smokers. *Inter J. Med. Sci.* 2013; 6(3): 29-34.
- Padmini E, Sundari BT. Erythrocyte Glutathione depletion impairs resistance to haemolysis in women consuming alcohol. *J Clin Biochem Nutr.* 2008; 42(1): 14-20.
- Blum J, Fridovich I. Inactivation of glutathione peroxidase by superoxide radical. *Arch Biochem Biophys.* 1985; 240(2): 500-8.
- Gul M, Kutay FZ, Temocin S, Hanninen O. Cellular and clinical implications of glutathione. *Ind J Exp Biol.* 2000; 38(7): 625-34.
- Raes M, Mihiels MC, Remacle J. Comparative study of the enzymatic defense systems against oxygen derived free radicals: the key role of glutathione peroxidase. *Free Radic Biol Med.* 1987; 3(1): 3-7.
- Lapenna D, Gioia S, Ciotani G, Mezzetti A, Uchino S. Glutathione related antioxidant defenses in human atherosclerotic plaques. *Circulation* 1998; 97(19): 1930-4.
- Zwirska KK, Jochem J, Rybus KB, Polaniak R, Birkner E. Assessment of blood superoxide dismutase, glutathione peroxidase activity and malondialdehyde concentration as oxidation status parameters in obese women. *Pol Arch Med Wewn.* 2003; 110(1): 725-31.