

A STOCHASTIC MODEL BY THE BIRTH AND DEATH PROCESS OF IRREDUCIBLE STATIONARY PROBABILITIES FOR THE GLP-1.

Dr. D. SargunaSundari

Assistant Professor Department of Mathematics AnnaiVailankanni Arts and Science College, Thanjavur, Tamilnadu, India.

Abstract: The Glucagon-like peptide-1 (GLP-1) inhibits appetite in part through regulation of soluble leptin receptors. GLP-1 is an incretin that it has the ability to decrease blood sugar levels in a glucose-dependent manner by enhancing the secretion of insulin. The time period of during weight loss maintenance, GLP-1 receptor agonist (GLP-1RA) administration may inhibits weight loss-induced increases in soluble leptin receptors there by preserving free leptin levels and preventing weight regain.

Here we develop a model for the customers arrive to the service according to a Poisson process of intensity λ and they immediately go to a free shop assistant, where they are received according to an exponential distribution of parameter μ .

$$p_k = \frac{1}{k!} e^{-\delta} \delta^k, k \in N_0$$

2000 Mathematics subject classification:

47N30,60H30,65C30

Index Terms: Birth-Death Processes, Stochastic Models, Glucagon-like peptide-1, leptin receptors.

I. INTRODUCTION

GLP-1 is a hormone secreted from the endocrine L cells in the intestine, which reduces food intake and when administered to obese subjects, induces weights loss. (Astrup, A., Rössner, S., ,2009). It is well established that GLP-1 inhibits food intake by inhibiting appetite. However, the leptin receptor may be involved in the appetite inhibition by GLP-1. (Iepsen, E. W., 2015) Many obese people have a history of several successful weight losses, but very few are able to maintain the weight loss over a longer period of time. After weight loss, several physiological mechanisms favoring weight regain are activated, including declining energy expenditure in combination with hunger and reduced satiety during eating, (Rosenbaum, M.,2008) Here it has been hypothesized that the reduction in free circulating leptin after weight loss is responsible for the weight regain and experiments have shown that leptin repletion after weight loss can reduce weight regain by decreasing energy intake.

II. NOTATIONS

$\frac{1}{\lambda}$ – mean inter arrival time

$\frac{1}{\mu}$ – mean service time

λ_k – Birth intensity at state E_k

μ_k – Death intensity at state E_k

III. BIRTH AND DEATH PROCESSES

Let $\{x(t) | t \in [0, D]\}$ be a stochastic process, which can be in the states E_0, E_1, E_2, \dots . The process can only move from one state to a neighboring state in the following sense. If the process is in state E_k , and we receive a positive signal then the process is transferred to E_{k+1} , and if instead we receive a negative signal (and $k \in N$), then the process is transferred to E_{k-1} . (Lawler, G. F. 2018)

We assume that there are non-negative constants λ_k and μ_k , such that for $k \in N$,

$$\begin{aligned} 1) \quad P\{\text{one positive signal is }] t, t + h[| x(t) = k\} \\ = \lambda_k h + \lambda \in (h). \end{aligned}$$

- 2) $P\{\text{one negative signal is }]t, t + h[\mid x(t) = k\} = \mu_k h + h \in (h).$
- 3) $P\{\text{no signal is }]t, t + h[\mid x(t) = k\} = 1 - (\lambda_k + \mu_k)h + h \in (h),$

λ_k the birth intensity at state E_k , and μ_k is called the death intensity at state E_k , and the process itself is called a birth and death process. If in particular all $\mu_k = 0$, if it is a birth process, and a death process, if all $\lambda_k = 0$.(Mejlbro, L. 2009)(Medhi, J. 2002)

A simple analysis shows for $k \in N$ and $h > 0$ that the event $\{X(t + h) = k\}$ is realized in one of the following ways:

- $X(t) = k$, and no signal in $]t, t + h[$.
- $X(t) = k - 1$, and one positive signal in $]t, t + h[$.
- $X(t) = k + 1$, and one negative signal in $]t, t + h[$.
- More signals in $]t, t + h[$.

We put $P_k(t) = p\{X(t) = k\}$ (3.1)

By a rearrangement and taking the limit $h \rightarrow 0$, we easily derive the differential equation of the process,

$$\begin{aligned} p'_0(t) &= -\lambda_0 p_0(t) + \mu_1 p_1(t) \text{ for } k = 0 \\ p'_k(t) &= -(\lambda_k + \mu_k) p_k(t) + \lambda_{k-1} p_{k-1}(t) + \mu_{k+1} p_{k+1}(t), \text{ for } k \in N \end{aligned} \tag{3.2}$$

In a special case of a pure birth process, where all $\mu_k = 0$, this systems is reduced to

$$\begin{aligned} p'_0(t) &= -\lambda_0 p_0(t) \text{ for } k = 0 \\ p'_k(t) &= -\lambda_k p_k(t) + \lambda_{k-1} p_{k-1}(t), \text{ for } k \in N \end{aligned} \tag{3.3}$$

If all $\lambda_k > 0$, we get the following iteration formula of the complete solution,

$$\begin{aligned} p_0(t) &= C_0 e^{-\lambda_0 t}, \text{ for } k = 0 \\ p_k(t) &= \lambda_{k-1} e^{-\lambda_k t} \int_0^t e^{\lambda_k T} p_{k-1}(T) dT + C_k e^{-\lambda_k t} \text{ for } K \in N. \end{aligned} \tag{3.4}$$

Let $\{X(t), t \in [0, +\infty]\}$ be a birth and death process, where all λ_k and μ_k are positive, with the exception of μ_0 and $\lambda_N = 0$, if there is a final state E_N . The process can be in any of the states, therefore, in analogy with the markov chain, such a birth and death process is called irreducible.(H, Constantin, 2016)

In Queuing theory, if there exists a state E_k , in which $\lambda_k = \mu_k$, then E_k is an absorbing state, because it is not possible to move away from E_k .

For the most common birth and death processes there exist non-negative constants p_k , such that $p_k(t) \rightarrow p_k$ and

$$p'_k(t) \rightarrow 0, \text{ for } t \rightarrow \infty.$$

These constants fulfill the infinite system of equation,

$$\mu_{k+1} p_{k+1} = \lambda_k p_k, \text{ for } k \in N_0,$$

which sometimes can be used to find the p_k .

If there is a solution (p_k) , which satisfies,

$$\begin{aligned} p_k &\geq 0 \text{ for all } k \in N_0, \\ \text{and } \sum_{k=0}^{\infty} p_k &= 1 \end{aligned}$$

The solution (p_k) is a stationary distribution, and the p_k are called the stationary probabilities.

In this case we have

$$p_k(t) \rightarrow p_k \text{ for } t \rightarrow \infty.$$

If $\{X(t) | t \in [0, \infty[$ is an irreducible process, then

$$p_k = \frac{\lambda_{k-1}\lambda_{k-2}\dots\lambda_1\lambda_0}{\mu_k\mu_{k-1}\dots\mu_2\mu_1} \cdot p_0 \tag{3.5}$$

where all $a_k > 0$.

The condition of the existence of a stationary distribution is then reduced to that the series $\sum_k a_k$ is convergent of finite sum $a > 0$.

In this case we have $P_0 = \frac{1}{a}$

IV. QUEUING PROCESSES

In the $M|M|1$ model is a poisson-input, exponential service, single server queue. The density functions for the inter arrival and service times are given.(Gross, D. 2008)

$$a(t) = \lambda e^{-\lambda t}$$

$$b(t) = \mu e^{-\mu t} ,$$

where $1/\lambda$ is the mean inter arrival time and $1/\mu$ is the mean service time.

Both the inter arrival and service time are exponential, and the arrival and conditional service rates are poisson, which gets us

$$p\{an \text{ arrival occurs in an interval of length } \Delta t\} = \lambda\Delta t + o(\Delta t)$$

$$p\{more \text{ than one arrival occurs in } \Delta t\} = o(\Delta t)$$

$$p\{a \text{ service completion in } \Delta t \text{ given the system is not empty}\} = \mu\Delta t + o(\Delta t)$$

$$p\{more \text{ than one service completion in } \Delta t \text{ given more than one in the system}\} = o(\Delta t)$$

The $M|M|1$ model is a simple birth-death process with $\lambda_n = \lambda$ and $\mu_n = \mu$ for all n .

Arrivals are ‘births’ to the system, since if the system is in state n (the state refers to the numbers of customers in the system), an arrival increases it to state $n + 1$.

Similarly, departures are death moving from state n to state $n - 1$.

Hence, the steady state equation are found to be

$$0 = -(\lambda + \mu)p_n + \mu p_{n+1} + \lambda p_{n-1}.$$

$$0 = -\lambda p_0 + \mu p_1$$

(or)

$$p_{n+1} = \frac{\lambda + \mu}{\mu} p_n - \frac{\lambda}{\mu} p_{n-1}$$

$$p_1 = \frac{\lambda}{\mu} p_0, \text{ which is a set of second order difference equation with constant coefficient.}$$

To solve the steady state difference equation for $\{p_n\}$ using the iterative method since the $M|M|1$ systems is a birth-death process with constant birth and death rates, we have

$$p_n = p_0 \prod_{i=1}^n \frac{\lambda}{\mu}$$

$$= p_0 \left(\frac{\lambda}{\mu}\right)^n$$

To find p_0 , we remember that probabilities must sum to 1 and it follows that

$$1 = \sum_{n=0}^{\infty} \left(\frac{\lambda}{\mu}\right)^n p_0$$

If $\rho = \frac{\lambda}{\mu}$ as the traffic intensity for the single-server queues.

$$p_0 = \frac{1}{\sum_{n=0}^{\infty} \rho^n}$$

Now $\sum_{n=0}^{\infty} \rho^n$ is an infinite geometric series which converges if and only if $\rho < 1$

Thus, the only way for a steady-state solution to exist, if and only if $\rho < 1$

Since, we know the sum of the terms of a convergent geometric series

$$\sum_{n=0}^{\infty} \rho^n = \frac{1}{1-\rho} (\rho < 1)$$

We find $p_0 = 1 - \rho$, which confirms the general results for p_0 we derived in the $G|G|1$.

Thus the full steady-state solution for the $M|M|1$ systems in the geometric probability function.

$$p_n = (1 - \rho)\rho^n \left(\rho = \frac{\lambda}{\mu} < 1\right)$$

The stochastic process is described by the birth and death process,

$$\{\alpha(t)/t \in [0, +\infty] \text{ and } \lambda_k = \lambda \text{ and } \mu_k = k\mu \text{ for all } k.$$

The process is irreducible, and the differential equation of the system are given by

$$p'_0(t) = -\lambda p_0(t) + \mu p_1(t) \text{ for } k=0$$

$$p'_k(t) = -(\lambda + k\mu)p_k(t) + \lambda p_{k-1}(t) + (k + 1)\mu p_{k+1}(t) \text{ for } k \in N \tag{4.1}$$

The stationary probabilities exist and satisfy the equations

$$(k + 1)\mu p_{k+1} = \lambda p_k, k \in N_0 \text{ of the solutions}$$

$$p_k = \frac{1}{k!} (\delta)^k e^{-\delta}, k \in N_0 \tag{4.2}$$

These are the probabilities that there are k customers in the system, when we have obtained equilibrium. The results of the Fig.2 are applied in the Eq. 4.2 are shown in Fig. 1.(Kumar, P. S., &Sundari, D. S. .2014)(Senthilkumar, P., &Sundari, D. S. 2015)




Figure 1 Endogenous GLP-1 in Control and Liraglutide by irreducible stationary probabilities.

V. EXAMPLE

Each study participant underwent three meal tests, at screening, after 8 weeks on very low calorie powder diet, and after 52 weeks with restrictive weight maintenance diet and recommendations (Astrup, A., Rössner, S., 2009). Various blood samples for measurements of Endogenous GLP-1 were taken before the intake of Fresutrin energy drink and every 15 min until 180 min after meal ingestion. (Kissileff, H. R., 2008) The 12% weight loss was successfully maintained in both the groups with no significant difference in weight between the beginnings of the weight loss maintenance program to after 52 weeks. In endogenous GLP-1 levels were significantly lower in the GLP-1RA group compared with the control group.




Figure 2 – Endogenous GLP-1 in Control and Liraglutide

VI. CONCLUSION

The mathematical model also reflects the same effects in the endogenous GLP-1, levels were lower in the GLP-1RA group compared with the control groups (fig 2) which are beautifully fitted with birth and death processes of irreducible stationary probabilities fig (1). The results matching with the mathematical and medical report.

REFERENCES:

[1] Astrup, A., Rössner, S., Van Gaal, L., Rissanen, A., Niskanen, L., Al Hakim, M., ... & NN8022-1807 Study Group. (2009). Effects of liraglutide in the treatment of obesity: a randomised, double-blind, placebo-controlled study. *The Lancet*, 374(9701), 1606-1616.

[2] Gross, D. (2008). *Fundamentals of queueing theory*. John Wiley & Sons.

[3] H, Constantin. (2016) *Markov Chains and Queuing Theory*. doi: 10.1007/978-1-4614-4084-0_5

[4] Iepsen, E. W., Lundgren, J., Dirksen, C., Jensen, J. E., Pedersen, O., Hansen, T., ... & Torekov, S. S. (2015). Treatment with a GLP-1 receptor agonist diminishes the decrease in free plasma leptin during maintenance of weight loss. *International Journal of Obesity*, 39(5), 834.

- [5] Kissileff, H. R., Thornton, J. C., Torres, M. I., Pavlovich, K., Mayer, L. S., Kalari, V., ... & Rosenbaum, M. (2012). Leptin reverses declines in satiation in weight-reduced obese humans. *The American journal of clinical nutrition*, 95(2), 309-317.
- [6] Kumar, P. S., & Sundari, D. S. (2014). A stochastic vasicek model for the glucose-induced glucagon-like peptide 1 secretion is deficient in patients with non-alcoholic fatty liver disease. *Arya Bhatta Journal of Mathematics and Informatics*, 6(2), 305-308.
- [7] Lawler, G. F. (2018). *Introduction to stochastic processes*. Chapman and Hall/CRC.
- [8] Medhi, J. (2002). *Stochastic models in queueing theory*. Elsevier.
- [9] Mejlbro, L. (2009). *Stochastic Processes 1*. Bookboon.
- [10] Rosenbaum, M., Sy, M., Pavlovich, K., Leibel, R. L., & Hirsch, J. (2008). Leptin reverses weight loss-induced changes in regional neural activity responses to visual food stimuli. *The Journal of clinical investigation*, 118(7), 2583-2591.
- [11] Senthilkumar, P., & Sundari, D. S. (2015) An euler-maruyama method to the black scholesde to the preserved incretia activity of glp-1 with type-2 diabetes patients. *Bulletin of Mathematics and Statistics Research*. 3(3). 118-123.