ISSN: 0975-3583, 0976-2833 VOL 12, ISSUE 04, 2021

Original Research Article

A Prospective Study to Evaluate the Pattern of Cervical PAP Smear Cytology and its Correlation with Clinical Findings in Tertiary Care Centre of North Bihar

Dr. Ranjan Kumar Rajan¹, Dr. MD Shakir Ahmad², Dr. Chittaranjan Dutta³

¹Assistant Professor, Department of Pathology, Darbhanga Medical College, Laheriasarai, Darbhanga, Bihar, India

²Assistant Professor, Department of Pathology, Darbhanga Medical College, Laheriasarai, Darbhanga, Bihar, India

³Associate Professor & Head, Department of Pathology, Darbhanga Medical College, Laheriasarai, Darbhanga, Bihar, India

Corresponding Author: Dr. MD Shakir Ahmad

Received :11-08-2021. Revised:21-08-2021. Accepted:20-09-2021

Abstract

Aim: the aim of the study to evaluate the pattern of cervical Pap smear cytology at a tertiary hospital and to correlate it with clinical findings.

Methods: The prospective study was carried out at Department of Pathology, Darbhanga Medical College, Laheriasarai, Darbhanga, Bihar, India for 1 year. The cellular material obtained on the spatula and cyto brush was quickly smeared on a clean glass slide, labeled, fixed in 95% ethyl alcohol immediately and subsequently stained by Pap stain. After staining, slides were mounted with DPX (Dibutyl phthalate Polystyrene xylene), screened and reported by cytopathologist under light microscope.

Results: Total 1000 cervico-vaginal smears studied during study period on patients, ranging from 18 to 63 years. Cytological findings broadly classified into unsatisfactory smears, normal and abnormal smears. There were 779(77.90%) abnormal Pap smears which include benign cellular changes of inflammation as well as Epithelial Cell Abnormalities (ECA), with 201(20.10) normal cases and 20(2%) unsatisfactory or inadequate samples. The age range of patients with epithelial cell abnormality was 22 to 63 years and the mean age was 44.1 ± 10.7 years. Total 440 (44%) showed inflammatory lesion, 12 (1.20%) showed atrophy, 8(0.80%) showed ASCUS, 5 (0.5%) showed ASC-H, 5(0.5%) showed HSIL, 10(0.10%) showed SCC, 2 (0.20%) AGUS, 110 (11%) showed metaplasia, 14 (0.14%) had radiation changes.

Conclusion:By proper implementation of Pap screening program, incidence of invasive cervical malignancy can be prevented due to early detection of cervical premalignant lesions. **Keywords:**Pap screening, metaplasia, cervical lesion

ISSN: 0975-3583, 0976-2833 VOL 12, ISSUE 04, 2021

Introduction

Cervical cancer is the fourth most common cancer worldwide. It is the second most common cancer after breast cancer in India with 96,922 new cases and 60,018 deaths in 2018.¹ Female in developing countries generally visit to gynecology department only when symptomatic with pain abdomen, discharge per vaginum, or menstrual abnormalities. Rarely, they come for routine screening for the detection of cervical abnormalities.² The Papanicolaou (Pap) smear was introduced in 1941 and became the standard screening test for cervical cancer and premalignant lesions.³ Cervical cancer is the most common cancer among women after breast and colorectal cancer in the world, but in developing country like India it is the leading cause of mortality and morbidity. Women in these countries usually present to the clinic only when they have symptoms, such as pain, discharge, and/or abnormal bleeding. Nearly 4 lacks new cases of cervical cancers are diagnosed annually worldwide and 80% of them are diagnosed in the developing countries.⁴ Cervical cancers can be prevented through early detection by means of effective screening techniques. Cervical Pap smear is a sensitive test for early screening of the cervical lesion. Though Pap smear is just a routine screening test, the overall sensitivity in detection of premalignant lesions like high grade squamous intraepithelial lesion (HSIL) is 70-80% and has been proved very effective in differentiating between inflammatory, premalignant and malignant lesions.⁵ Thus the epithelial changes can be treated, preventing the cervical cancer.⁶, In 1988, the Bethesda system of terminology has been introduced to sub-classify the lesions into high grade and low grade squamous intraepithelial lesions (SIL) for Pap smear reporting and some studies reported comparison of various terminologies.^{7,8} The present study is intended to evaluate the pattern of cervical Pap smear cytology at a tertiary hospital and to correlate it with clinical findings.

Material and methods

The prospective study was carried out at Department of Pathology, Darbhanga Medical College, Laheriasarai, Darbhanga, Bihar, India for 1 year.

Methodology

Total 1000 patients were screened. The patients were in the age range of 18-63 years, having complaints like vaginal discharge, bleeding per vagina or something coming out per vagina, post-coital bleeding, intermenstrual bleeding, and pain in lower abdomen. History and symptoms along with parity were recorded. Smears were taken by trained technician using modified Ayres wooden spatula which was inserted and rotated 360 over cervix. Both ectocervix and endocervix were sampled. The cellular material obtained on the spatula and cyto brush was quickly smeared on a clean glass slide, labeled, fixed in 95% ethyl alcohol immediately and subsequently stained by Pap stain. After staining, slides were mounted with DPX (Dibutyl phthalate Polystyrene xylene), screened and reported by cytopathologist under light microscope according to the 2014 Bethesda system.

Results

Total 1000 cervico-vaginal smears studied during study period on patients, ranging from 18 to 63 years. Cytological findings broadly classified into unsatisfactory smears, normal and abnormal smears. There were 779(77.90%) abnormal Pap smears (benign cellular changes of inflammation as well as Epithelial Cell Abnormalities (ECA), with 201(20.10) normal cases and 20(2%) unsatisfactory or inadequate samples.

ISSN: 0975-3583, 0976-2833 VOL 12, ISSUE 04, 2021

The age range of patients with epithelial cell abnormality was 22 to 63 years and the mean age was 44.1 ± 10.7 years.

Total 476 (47.6%) showed inflammatory lesion, 12 (1.20%) showed atrophy, 8(0.80%) showed ASCUS, 5 (0.5%) showed ASC-H, 5(0.5%) showed HSIL, 1(0.10%) showed SCC, 2 (0.20%) AGUS, 110 (11%) showed metaplasia, 14 (0.14%) had radiation changes.

Table 1. Age wise distribution of total number of patients				
Age	No. of patients	Percentage		
Below 30	18	1.80		
30-40	131	13.10		
40-50	341	34.10		
50-60	298	29.80		
Above 60	212	21.20		

Table 1: Age wise distribution of total number of patients

Diagnosis			No. of cases	Percentage
Unsatifactory Smear		20	2%	
	Normal		201	20.10%
NILM		Non-specific	440	44%
	Inflammatory	Candida	29	2.90%
		Trichomonas	7	0.70%
	Reactive		146	14.6%
	Metaplasia		110	11%
Atrophy			12	1.20%
Radiation			14	1.40%
ASCUS			8	.80%
ASC-H		5	0.5%	
HSIL			5	0.5%
SCC			1	0.10%
AGUS			2	0.20%

Table 2: Cytological findings

Discussion

Cancers of uterine cervix and breast are the leading malignancies seen in females of India. There should be an effective mass screening program aimed at specific age group for detecting precancerous condition before they progress to invasive cancers.^{9,10} Cervical cytology is currently widely used as the most effective cancer screening modality. Objective data from hospital-based studies are required in order to detect the efficiency of the screening test. This study contributes to assessing current levels of cervical screening in the tertiary teaching hospital in bihar, India. In our study, the mean age of patients with abnormal smears was 44.1 ± 10.7 years. Similar finding was detected by other studies.¹¹ This study determines 201cases (20.10%) of normal findings, inflammatory lesions in 476 cases (47.6%). The Epithelial Cell Abnormality (ECA) rate, that is the total of ASCUS, ASC-H, LSIL, HSIL, AGUS and carcinoma is 2.1 %.^{12,13}

Our study revealed ASCUS (0.80%) to be the most common epithelial cell abnormality, most of found in age group of 30-50 years of age. Similar results were obtained in other studies which also concluded ASCUS to be the most common epithelial cell abnormality.^{13,14} ASCUS progresses to LSIL, HSIL and SCC. AGUS progresses to adenocarcinoma.^{14,15}

ISSN: 0975-3583, 0976-2833 VOL 12, ISSUE 04, 2021

Edelman et al, studied Pap smears from 29295 females over a period of one year and the Pap smear abnormalities were as follows: 9.9% ASC-US, 2.5% LSIL, 0.6% HSIL, and 0.2% invasive cancer.¹⁶ Study by Banik U revealed the following scenario: 0.18% ASCUS, 0.12% Atypical glandular cells (AGC), 6.36% LSIL, 1.18% HSIL and 0.35% malignancy.

In our study shows ASCUS 0.80%, ASC-H 0.5%, HSIL 0.5%, SCC 0.10% and AGUS 0.20%. One of the significant discrepancies between our study and the previously published data from other countries is the higher rate of ASC-US and lower rate of LSIL. We assume that as the women included in our study were routinely screened and/or re-screened, they presented with an early form of cytological interpretation in the cervical smear, and thus, ASCUS rate was higher. Most common age to develop carcinoma cervix is between 40 and 50 years and the precursor lesions occur 5-10 years prior to developing invasive cancer. Various screening test for cervical cancer like Pap smear, liquid Pap cytology, automated cervical screening techniques, visual inspection of cervix after Lugol's Iodine and acetic acid application, speculoscopy, cervicography should be started for early detection of premalignant lesions.

Conclusion

This study emphasized the importance of Pap smears screening for early detection of premalignant and malignant lesions of cervix. Larger studies are required to estimate the pattern of cervical cytological abnormalities along with detection of common HPV strains in cervical cancer in Indian population. Proper implementation of Pap screening program, incidence of invasive cervical malignancy can be prevented due to early detection of cervical premalignant lesions.

Reference

- 1. Available from: http: //www.cancerindia. org.in/globocan- 2018-india-factsheet. [Last accessed on 2019 Sep 26].
- 2. Saslow S, Solomon D, Lawson HW, Killackey M, Kulasingam SL, Cain J, *et al.* American cancer society, american society for colposcopy and cervical pathology, and American society for clinical pathology screening guidelines for the prevention and early detection of cervical cancer. J Low Genit Tract Dis 2012;16:175-204.
- 3. Papanicolaou GN, Traut HF. The diagnostic value of vaginal smears in carcinoma of the uterus. Am Jr Obstet Gynecol. 1941;42:193-205.
- 4. Patel MM, Pandya AN, Modi J. Cervical pap smear study and its utility in cancer screening, to specify the strategy for cervical cancer control. National Journal of Community Medicine.2011;2(1).
- 5. Kazi BM, Karamat KA. Pap smear screening for Precancerous conditions of the cervical cancers. Pak J Med. Res.2005;44(3):111-3
- 6. Maryem A, Ghazala M, Arif HA, Tamkin K. Smear pattern and spectrum of premalignant and malignant cervical epithelial lesions in postmenopausal Indian women: a hospital based. Study. Diagnostic Cytopathology; 40(11):976-983, 2011.
- 7. Jonathan S.B. Berek and Novak's Gynaecology. 14th ed. Philadelphia: Lippincott William Wilkins. 2006;569-75.
- 8. Leopold K. The new bethesda system for reporting results of smears of uterine cervix. 10. Journal of National Cancer Institute. 1990;82(12):988-90.
- 9. Kalkar RA, Kulkarini Y. Screening for cervical cancer: an overview. Obstet Gynecol India. 2006;56(2):115-22.
- 10. Khan MS, Raja FY, Ishfaq G, Tahir F, SubhanF,Kazi BM et al. Pap smear Screening for Precancerous conditions of the cervical cancers. Pak J. Med. Res. 2005;44(3):111-3.

ISSN: 0975-3583, 0976-2833 VOL 12, ISSUE 04, 2021

- 11. Turkish Cervical Cancer And Cervical Cytology Research Group: Prevalence of cervical cytological abnormalities in Turkey. Int J Gynaecol Obstet. 2009;106:206-9.
- 12. Ghaith JE, Rizwana BS. Rate of Opportunistic Pap smear Screening and Patterns of Epithelial Cell Abnormalities in Pap Smears in Ajman, United Arab Emirates. Sultan Qaboos Univ Med J. 2012;12(4):473-8.
- 13. Saslow D, Solomon D, Lawson HW, Killackey M, Kulasingam S, Cain et al. American Cancer Society, American Society for Colposcopy and Cervical Pathology, and American Society for Clinical Pathology Screening Guidelines for the Prevention and Early Detection of Cervical Cancer. Journal of Lower Genital Tract Disease. 2012;16(3):175-204.
- 14. Edelman M, Fox A. Cervical Papanicolau smear abnormalities in inner Bronx adolescents: Prevalence, progression, and immune modifiers. Cancer (cancer cytopathology). 1999;87:184-9.
- 15. Khan MS, Raja FY, Ishfaq G, Tahir F, SubhanF,Kazi BM et al. Pap smear Screening for Precancerous conditions of the cervical cancers. Pak J. Med. Res. 2005;44(3):111-3.
- 16. Banik U, Bhattacharjee P, Ahamad SU, Rahman Z. Pattern of epithelial cell abnormality in Pap smear: A clinicopathological and demographic correlation. Cyto Journal. 2011;8:8.