

**ORIGINAL RESEARCH****A Study on Relative Contribution of Hardiness, Self-esteem and Learned Helplessness in Relation to Alcohol Use****<sup>1</sup>Lovesampuranjot Kaur, <sup>2</sup>Bhupinder Kaur**<sup>1</sup>Professor, <sup>2</sup>Assistant Professor, Faculty of Nursing, Desh Bhagat University, Mandi Gobindgarh, Punjab, India**Correspondence:**

Lovesampuranjot Kaur

Professor, Faculty of Nursing, Desh Bhagat University, Mandi Gobindgarh, Punjab, India

**Abstract**

The aim of the present study was to determine the relative contribution of Hardiness, Self-esteem and Learned Helplessness in relation to alcohol use. A sample of 300 alcoholics (adult males) was selected from community and de-addiction centres from Punjab. The measures used were Short Alcohol Dependents Data Questionnaire (SADD, Raistrick, Dunbar and Davidson 1983), Hardiness Scale (Kobasa & Kahn, 1982), Self-esteem Inventories Adult Form (Coopersmith's, 1981) and Learned Helplessness Scale by Dhar, U., Kohli, S, & Dhar, S. (1987). Findings indicated that Hardiness, Self-esteem and Learned Helplessness contributed significantly for social drinkers whereas did not contribute significantly to increase level of alcohol use for problem drinkers and alcohol dependents.

**Key words:** Hardiness, self-esteem, learned helplessness, alcoholism.

Alcoholism and drug abuse is widely recognized as a serious problem world over with severe psychological, social and physical consequences. Hence the problem of drug and alcohol abuse is not unique either to India or to present times (Sachene, 1990) but is a chronic menace.

The health care system is greatly affected by alcoholism. In India, 10% of adults entering private physician's clinics are alcoholics and 15-40% of adult admissions to general hospitals are for alcohol related problems. (W.H.A. Report, 2002). One fact comes to the forefront while analyzing the whole scenario that is what makes certain drinkers strictly remain social drinkers while others further deteriorate to drinking as a habit and become addicts/dependents.

**DSM IV-TR** (A.P.A., 2005) differentiates different levels of alcohol use as follows: -

**Alcohol dependence** is characterized by at least three of specific signs or symptoms from inability to control the amount consumed interferences with work, school or social activities, tolerance, withdrawal and duration of problem being at least for a month.

**Problem drinkers**

These are people who can not drink in a controlled manner, or people whose drinking at one time has adversely affected their health or caused them any economic, professional, legal or personal problems (National Institute on Alcohol Abuse & Alcoholism, 1992).

**Social drinking**

Can be defined as drinking pattern that is found to be acceptable to the society in which they occurs on an infrequent basis during social occasions that may call for alcohol to be present

and/or consumed. Those individuals who engage in social drinking generally only have one or two drinks and are easily able to stop drinking at that time. Social drinking is defined as such because under normal circumstances, the individual would probably not choose to consume alcohol but may do so only due to the social situation (United State Department of Health and Human Services, 1992).

Research suggests that certain personality factors/traits may play an important role in both the development and maintenance of alcohol dependence (Barnes, 1980). Characteristics that have been identified include impulsivity, negative self concept, weak ego, low social conformity, neuroticism and introversion. It has also been associated with antisocial personality and depressive response styles (Leigh, 1985). This may be explained by the inability of an individual with antisocial personality to anticipate the aversive consequences of his or her behavior. It is likely that in an effort on the part of that person to manage negative emotional evaluations, he may indulge in substance abuse. Further it may be an impulsive act towards anxiety relief. Achievement of relief then provides the positive reinforcement to continue abusing the substance.

### **Hardiness**

The term hardiness was introduced by Kobasa (1979) to refer to the personality style which keeps the person healthy even after prolonged exposure to stress. Hardy people are hypothesized to possess three general characteristics: commitment, control & challenge.

### **Commitment**

Hardy people show deeper involvement in whatever they do and have a tendency to perceive these activities as worth doing. Persons strong in commitment have a strong sense of purpose and direction and do not easily give up under pressure. Commitment is reflected in the ability to feel activity involved with others and a belief in the truth, value and importance of one's self and one's experience (Huang & Wagnitd, 1995; Tartasky, 1993). Adverse situations are ultimately seen as meaningful and interesting (Maddi & Kobasa, 1985).

### **Control**

They have a tendency to feel and act in an influential manner in the face of varied contingencies of life. They feel both capable and empowered to achieve desired outcomes (Kobasa, 1979). They act as they are influential in contingencies of life, events are perceived as a natural outgrowth to the individuals actions and not as unexpected experiences (Kobasa et al., 1982).

### **Challenge**

Hardy people tend to perceive changes as a challenge, for them anticipation of changes are interesting incentives to growth rather than threat to security. Challenge reflects the belief that change is not a threat to personal security, but an opportunity for personal development and growth (Kobasa & Maddi, 1984). Hardiness reduces unhealthy effects of stress in two ways: (1) it improves health by acting as a buffer to stressful life events (Kobasa & Puccetti, 1983) and (2) it directly reduces the strain by decreasing the use of unsuccessful coping strategies (Kobasa et al., 1982).

### **Self-esteem**

Self-esteem refers to an individual's sense of his or her value or worth, or the extent to which a person values, approves of, appreciates, prizes or likes him or herself. The most broad and frequently cited definition of self esteem is by Rosenberg (1965), who described it as a favorable or unfavorable attitude towards the self. Self esteem is generally considered the

evaluative component of the self concept, a broader representation of the self that includes cognitive and behavioral aspects as well as evaluative or affective ones. While the construct is most often used to refer to a global sense of self worth, narrower concepts such as self confidence or body esteem are used to imply a sense of self esteem in more specific domains. It is also widely assumed that self esteem functions as a trait, that is, it is stable across time within individuals (Blascovich & Tomaka, 1991).

The term learned helplessness describes an organism's reaction when it is faced with important events that cannot be altered by its voluntary responses. Learned helplessness is both a behavioral state and a personality trait of one who believes that control has been lost over the reinforcers in the environment. These negative expectations lead to helplessness, passivity and an inability to assert oneself.

### **Objective**

To determine relative contribution of hardiness, self esteem and learned helplessness on different levels of alcohol use.

### **Hypothesis**

Hardiness, self-esteem & learned helplessness will significantly contribute to increase level of alcohol use.

### **Tools**

The tools were selected in accordance with the aims and objectives of the study. While selecting the tools, psychometric properties, nature of sample, competence of the investigator in scoring and interpretation was taken into consideration. The scales had to be adapted/translated for the sample of the present study.

The Tools used for the study were as follows:

1. Short Alcohol Dependence Data Questionnaire (Raistrick, Dunbar, & Davidson, 1983).
2. Hardiness Scale (Kobasa & Kahn, 1982).
3. Self-esteem Inventories Adult Form (Coopersmith's, 1981).
4. Learned Helplessness Scale (Dhar, U., Kohli, S, & Dhar, S. 1987).

**1. Short Alcohol Dependence Data questionnaire (SADD; Raistrick, Dunbar, & Davidson, 1983).** This measure is a 15-item measure that assesses the range of current state alcohol dependence (i.e., behavioral, subjective, and psychobiological changes associated with alcohol dependence).

**2. Hardiness Scale:** To measure the hardiness level of subjects Psychological Hardiness Scale (Kobasa & Kahn, 1982) was used. The scale consists of 12 items positively and negatively keyed covering the important dimensions of hardiness as commitment, control and challenge. The scale was administered to the subjects after translating into Punjabi. Scoring was done in accordance to the manual of the scale. The reliability coefficient of the translated scale was found to be 0.628 by the investigator. The validity of the scale was also found to be 0.543.

**3. Self-esteem Inventories- Adult Form,** The scale developed by Coopersmith's(1981) it is uni-dimensional scale which measures the self-esteem level. This form is used with persons aged 16 and above. It consists of 25 items which are to be answered "like me or unlike me". It has both positive and negative items to be answered. Maximum score is 100. High score corresponds to high self-esteem. The author reported its internal consistency reliability (determined by Kuder-Richerdson formula) 0.81 and 0.86. Test-retest reliability reported by author to be 0.88 and 0.70 respectively.

**4. Learned Helplessness Scale:** (Dhar, U., Kohli, S., & Dhar, S.,1987). To measure the learned helplessness of subjects the learned helplessness scale was used. This scale consists

of 15-items. All items have to be answered in positive, negative and uncertain, and that no statement is to be left out right item was scored as 3, wrong 1 and uncertain as 2.

### Sample

The final sample of the study consisted of 300 adult males. They were in the age group of 25-45 years. This sample was selected out of a larger sample of 500 subjects, so as to have equal numbers of Social Drinkers, Problem Drinkers and Alcohol Dependents belonging to rural and urban areas of Punjab.

### Result and Discussion

To check the causation of hardiness, self esteem and learned helplessness on different Levels of Alcohol Use stepwise regression analysis,  $R^2$  and R along with F were calculated and the values are given in the tables 1, 2 and 3 below.

**Table 1: Stepwise Regression Analysis for Social Drinkers**

Predictor Variable/s	Degree of Freedom	$R^2$	R	F
<b>Model I</b>				
Learned Helplessness	99	0.180	0.424	21.488*
Learned Helplessness + Self-esteem	98	0.226	0.476	14.186*
Learned Helplessness + Self-esteem + Hardiness	97	0.188	0.434	9.901*
<b>Model II</b>				
Self-esteem	99	0.002	0.039	0.153
Self-esteem + Hardiness	98	0.007	0.084	0.342
Learned Helplessness + Self-esteem + Hardiness	97	0.188	0.434	9.901*
<b>Model III</b>				
Hardiness	99	0.006	0.080	0.630
Learned Helplessness + Hardiness	98	0.197	0.444	11.927*
Learned Helplessness + Self-esteem + Hardiness	97	0.188	0.434	9.901*

\*Significant at 0.01 level

\*\* Significant at 0.05 level

Model I, II and III of Table 1 reveals that for Social Drinkers values of  $R^2$  of Learned Helplessness, Self-esteem and Hardiness are 0.180, 0.002 and 0.006 respectively. 18% of Alcohol use is thus predicted by Learned Helplessness, 0.2% by Self-esteem and 0.6% by Hardiness. The value of  $R^2$  for Learned Helplessness, Self-esteem and Hardiness taken together is 0.188, thus 18.8% of Alcohol use is predicted by Learned Helplessness, Self-esteem and Hardiness taken together. The F value for the conjoint  $R^2$  is 9.901 which is significant at 0.01 level of significance. This leads to the conclusion that Learned Helplessness, Self-esteem and Hardiness conjointly predict Alcohol use among Social Drinkers more as compared to their separate prediction.

**Table 2: Stepwise Regression Analysis for Problem Drinkers**

Predictor Variable/s	Degree of Freedom	$R^2$	R	F
<b>Model I</b>				
Learned Helplessness	99	0.051	0.226	5.255**

Learned Helplessness + Self-esteem	98	0.052	0.229	2.684
Learned Helplessness + Self-esteem + Hardiness	97	0.073	0.27	1.781
<b>Model II</b>				
Self-esteem	99	0.021	0.144	2.062
Self-esteem + Hardiness	98	0.021	0.144	1.027
Learned Helplessness + Self-esteem + Hardiness	97	0.073	0.27	1.781
<b>Model III</b>				
Hardiness	99	0.001	0.025	0.061
Learned Helplessness + Hardiness	98	0.051	0.226	2.605
Learned Helplessness + Self-esteem + Hardiness	97	0.073	0.27	1.781

\*Significant at 0.01 level

\*\* Significant at 0.05 level

Model I, II and III of Table 2 reveals that for Problem Drinkers values of R<sup>2</sup> of Learned Helplessness, Self-esteem and Hardiness are 0.051, 0.021 and 0.001 respectively. 5.1% of Alcohol use is thus predicted by Learned Helplessness, 2.1% by Self-esteem and 0.1% by Hardiness. The value of R<sup>2</sup> for Learned Helplessness, Self-esteem and Hardiness taken together is 0.073, thus 7.3% of Alcohol use is predicted by Learned Helplessness, Self-esteem and Hardiness taken together. The F value for the conjoint R<sup>2</sup> is 1.781 which is not significant. This leads to the conclusion that Learned Helplessness, Self-esteem and Hardiness conjointly do not predict Alcohol use among Problem Drinkers more as compared to their separate prediction.

**Table 3: Stepwise Regression Analysis for Alcohol Dependents**

Predictor Variable/s	Degree of Freedom	R <sup>2</sup>	R	F
<b>Model I</b>				
Learned Helplessness	99	0.008	0.090	0.807
Learned Helplessness + Self-esteem	98	0.018	0.133	0.872*
Learned Helplessness + Self-esteem + Hardiness	97	0.04	0.2	1.510
<b>Model II</b>				
Self-esteem	99	0.000	0.003	0.001
Self-esteem + Hardiness	98	0.033	0.182	1.653
Learned Helplessness + Self-esteem + Hardiness	97	0.04	0.2	1.510
<b>Model III</b>				
Hardiness	99	0.032	0.172	3.234
Learned Helplessness + Hardiness	98	0.035	0.187	1.759
Learned Helplessness + Self-esteem + Hardiness	97	0.04	0.2	1.510

\*Significant at 0.01 level

\*\* Significant at 0.05 level

Model I, II and III of Table 3 reveals that for Alcohol Dependents values of R<sup>2</sup> of Learned Helplessness, Self-esteem and Hardiness are 0.008, 0.000 and 0.032 respectively. 0.8% of

Alcohol use is thus predicted by Learned Helplessness, 0.000% by Self-esteem and 3.2% by Hardiness. The value of  $R^2$  for Learned Helplessness, Self-esteem and Hardiness taken together is 0.04, thus 4% of Alcohol use is predicted by Learned Helplessness, Self-esteem and Hardiness taken together. The F value for the conjoint  $R^2$  is 1.510 which is not significant. This leads to the conclusion that Learned Helplessness, Self-esteem and Hardiness conjointly do not predict alcohol use among Alcohol Dependents more as compared to their separate prediction.

On the basis of tables 1, 2 and 3 it can be concluded that hypothesis 8 which states that “Hardiness, self-esteem & learned helplessness will significantly contribute to Level of Alcohol Use,” is partially rejected. The hypothesis is accepted for Social Drinkers where as rejected for Problem Drinkers and Alcohol Dependents.

The result shows that for Problem Drinkers and Alcohol Dependents- Hardiness, Self-esteem & Learned Helplessness do not significantly contribute to Level of Alcohol Use, where as for Social Drinkers- Hardiness, Self-esteem & Learned Helplessness significantly contribute to Level of Alcohol Use.

### Conclusion

Based on the finding it was concluded that majority of the alcoholism are dependent on self esteem but less dependent on hardness and helplessness,

### References

1. Sachene, A. H.(1990). Objective and Subjective dimensions of family burden. *Social Psychiatry and Psychiatric Epidemiology*, 25(6) 289-297.
2. W.H.A. Report (2002). c.f. Caetano, R., Thomas, F. (2006). Diagnosis of alcohol dependence in epidemiological surveys: *an epidemic of youthful alcohol dependence*. *Addiction*, 101 (Supp. 1) 2006. 111-114.
3. National Institute of Alcohol Abuse and Alcoholism, (1992). When Do Social Drinkers Becomes Alcoholics. [pubs.niaaa.nih.gov](http://pubs.niaaa.nih.gov), No. 16,315.
4. U.S. Department of Health & Human Services (1992). c. f. Guidelines by the U.S Departmental of Agriculture and Human Services. No. 16, 316-317.
5. Barnes, G. E. (1980). Characteristics of the clinical alcoholic personality. *Journal of Studies on Alcohol*, 41, 894-910.
6. Leigh, G. (1985). Psychological factors in the etiology of Substance abuse. c. f. Bratter, T. E., & Forrest, G. G. *Alcoholism and Substance Abuse; Strategies for Clinical Intervention*. New York, The Free Press. 220-240.
7. Kobasa, S. C. (1979). Stressful life events, personality and health: an inquiry into hardiness. *Journal of Personality and Social Psychology*, 37, 1-11.
8. Huang, M., & Wagnitd, G., (1995). Another book at Hardiness. *Image*. 23, 257-259.
9. Maddi, S.R. & Kobasa, S. (1984). *The hardy executive: Health under stress*. Homewood, IL: Dow Jones-Irwin.
10. Kobasa, S. C. (1979). Stressful life events, personality and health: an inquiry into hardiness. *Journal of Personality and Social Psychology*, 37, 1-11.
11. Kobasa, S. C., Maddi, S. R., & Kahn, S. (1982). Hardiness and health: A prospective study. *Journal of Personality and Social Psychology*, 42: 168–177.
12. Kobasa, S., Maddi, S., Puccetti, M. & Zola, M. (1985). Effectiveness of hardiness, exercise, and social support as a resource against illness. *Journal of Psychometric Research*, 29 (5), 525-533.
13. Blascovich, J., & Tomaka, J. (1991). Measures of self-esteem. In J.P. Robinson, Shaver, P.R. & Wightsman, L.S. (Eds.) *Measures of personality and social psychological attitudes*, Volume I. San Deiego, CA: Academic Press.

14. Raistrick, D., & Davidson, R.(1983). Alcohol Dependence Data Questionnaire. *British Journal of Addiction*, 78, 89-95.
15. Kobasa, S. C., & Kahn, S. (1982). Hardiness and health: A prospective study. *Journal of Personality and Social Psychology*, 43: 196–202.
16. Coopersmith, S. (1959). A method for determining types of self esteem. *Journal of Abnormal Social Psychology*, 59, 87-94.
17. Dhar, U., Kohli, S., & Dhar, S. (1987). Manual for the Learned Helplessness Scale (LH-Scale). Bhiwani: Haryana Council of Psychological Researches.