

## Original Research Paper

# An Intrathecal butorphanol and fentanyl as an adjuvant to bupivacaine (heavy) for Abdominal hysterectomy (Comparative Study)

Dr. Abhay Raj Yadav<sup>1</sup> (Assistant professor), Dr. Priyanka Jaiswal<sup>2</sup> (Senior Resident),  
Dr. Anshu Priyanka Lakra<sup>3</sup> (Senior Resident), Dr. Mohan Babu Nema<sup>4</sup> (Assistant Professor)

<sup>12</sup>Department of Anaesthesiology, SSMC, Rewa

<sup>3</sup>Department of Anaesthesiology, Aiims, Bhopal, M.P.

<sup>4</sup>Department of Anaesthesiology, MGM Medical College, Indore, M.P.

Corresponding Author:- Dr. Mohan Babu Nema

## ABSTRACT

### INTRODUCTION

Neuraxial block for lower limb surgeries are becoming popular as it has many advantages over general anaesthesia. Spinal anaesthesia consists of temporary interruption of nerve transmission in the subarachnoid space produced by the injection of a local anaesthetic solution in the subarachnoid space. Neuraxial opioids are widely used in conjunction with local anesthetics (LA) as they permit the use of lower dose of LA while providing adequate anesthesia and analgesia.

The objective of the study was to compare the efficacy of butorphanol and Fentanyl as an adjuvant to local anaesthetics in relation to onset, degree and recovery time of sensory and motor blockade and Hemodynamic changes in orthopaedic procedures done under spinal anaesthesia.

**Materials & Methods** The present study were conducted in the Department of anaesthesiology, Shyam Shah Medical College and Gandhi memorial Hospital, Rewa on patients of diagnosis were abdominal hysterectomy. Surgeries scheduled for elective surgery.

A detailed physical examination was carried out a day before the proposed surgery, Written and informed consent before participation in study were taken.

**Inclusion Criteria:-** After obtaining approval from institutional ethics committee and informed written consent from patients. 80 consented adult patient of ASA Grade I & II, Aged between 30-60 years, were included in the study.

**Exclusion criteria:-** Patients refusal to participate in the study, contraindications for spinal anesthesia, Sensitivity to study drugs, Recent onset of MI.

**KEYWORDS:** Hyperbaric Bupivacaine, Fentanyl, Butorphanol, Subarachnoid Block, Post-Operative Analgesia

## 1. INTRODUCTION

Spinal anaesthesia has emerged as an important technique, with simplicity, effectiveness, safety and success[1]. To improve the effect and duration of spinal anesthesia various drugs are used as an adjuvant to hyperbaric bupivacaine. Butorphanol is a lipophilic opioid agonist-antagonist analgesic with a published affinity for opioid receptors in vitro of 1:4:25 (mu: delta: kappa). A dose dependent increase in the duration of analgesia provided by epidural butorphanol for relief of post-caesarean section pain[2&3]. Fentanyl, a highly lipid soluble,

pure  $\mu$ -agonist opioid with rapid onset and short duration of action, has been used with various local anesthetics for a wide variety of surgical procedures[4,5&6].

## 2. MATERIALS AND METHODS

This prospective, observational hospital based study was conducted after approval from the institutional ethics committee and written informed consent of patients. Eighty patients, aged 30- 60 years, belonging to ASA physical status I or II and scheduled for elective, abdominal hysterectomy surgeries were divided into two groups.

Group A: - Intrathecal injection of 25 $\mu$ g butorphanol with 15 mg of 0.5 % Bupivacaine (heavy).

Group B: -Intrathecal injection of 25 $\mu$ g fentanyl with 15 mg of 0.5 % Bupivacaine (heavy).

The total volume injected was 3.5 ml in both the groups. Baseline vitals like Heart rate, Non-invasive blood pressure, respiratory rate, ECG, oxygen saturation were recorded. All patients were preloaded with 10ml/kg ringer lactate. Patients were placed in sitting position on the operation table. With strict aseptic precautions, midline approach subarachnoid block was achieved in L3-L4 space with 25G disposable Quincke spinal needle. Patients were immediately placed in the supine position with no tilt given to the table. The onset of sensory analgesia was tested by pinprick ,checking in a caudal to cephalic direction. Time taken to achieve highest sensory level was noted. Time taken for onset of maximum motor blockade was noted. Maximum grade of motor blockade achieved using modied Bromage score was noted. Time to return of Modied Bromage score to zero was recorded. Routine monitoring of pulse, BP, SpO<sub>2</sub>, ECG was instituted intraoperatively. Fluid administration was continued intra-operatively. Decrease in mean arterial pressure greater than 15% below the pre-anaesthetic base line value was treated with incremental doses of injection Mephenteremine 6 mg IV. A decrease in heart rate below 50 beats/min was treated with incremental doses of atropine 0.3 mg IV. Post operatively, pain assessment was carried out by VAS and duration of motor block was assessed by Bromage scale. Intramuscular injection diclofenac (75 mg) was given in the gluteal region as rescue analgesic on demand. At that time, VAS score was recorded, duration of effective analgesia was measured as time from intrathecal drug administration to patients 1st request for analgesic either in recovery room or in ward. Patient was kept under observation for a total period of 24 hrs. to observe for the total number of doses of analgesic required and any side-effect .

## STATISTICAL ANALYSIS

Difference between the groups in the demographic data and baseline values were analyzed using unpaired t-test. Analysis was performed using software IBM SPSS statistics for windows, Data were presented as mean  $\pm$  standard deviation. A  $P < 0.05$  was considered statistically significant.

## 3. RESULT:

Both The groups were comparable with respect to age, sex, weight and ASA physical status.

**Table 1: Demographic profile of patients:**

Demographic profile	Butorphanol (GP-A) (Mean $\pm$ SD)	Fentanyl ( GP-B) (Mean $\pm$ SD)	p-value
Age(yrs)	31.06 $\pm$ 4.96	30.13 $\pm$ 5.34	>0.05
Weight(kg)	48.13	48.06	>0.05

**Table 2: Characteristics of Sensory block**

Parameters	Butorphanol (GP-A) Mean SD	Fentanyl (GPB) Mean SD	P-Value
Highest Sensory level (median range)	T7 (T 6-8)	T8 (T5 - 8)	-
Time from injection to highest sensory level (min)	7.0± 1.8	7.2 ± 2.1	P>0.05
Time of two segment regression (min)	106 ± 21	85 ± 25	P<0.001
Time for sensory regression to S2 (min)	158 ± 22	P<0.001	P<0.001

**Times are presented in minutes (mean ± SD) .P < 0.001**



The highest sensory level achieved and the times to reach peak sensory level were comparable among the two groups. Significantly slower two segment regression were observed in the group A receiving butorphanol as compared to fentanyl and time to S2 regression were significantly more with intrathecal butorphanol (158 ± 22min.) than intrathecal fentanyl (135.36 ± 22min.).

**Table No. 3: Characteristics of motor block**

There was no difference in the onset of grade III motor block in Butorphanol (A) and

Parameters	Butorphanol (GP-A) Mean SD	Fentanyl (GPB) Mean SD	P-Value
Onset of max. motor block (min)	8.2 ± 3.5	8.2 ± 3.5	P>0.05
Duration of grade III motor block (min)	110 ± 20	110 ± 20	P<0.001
Duration of grade II motor block (min)	140 ± 28	132 ± 25	P<0.001
Duration of grade I motor block (min)	165 ± 30	162 ± 32	P<0.001

Fentanyl (B) group (8.0 ± 2.8 ,8.2 ± 3.5 minutes), respectively, but the duration of grade III, II or I motor block was higher in Butorphanol (A) as compare Fentanyl (B) group.



**Table No.4: Comparison hemodynamic variables between two groups**

Parameters	Mean±SD values		P-Value
	GP-A	GP-B	
Systolic BP(mmHg)	114.28±6.28	115.28± 5.04	P>0.05
Diastolic BP(mmHg)	74.57± 4.32	75.42± 3.42	
Heart rate (/min.)	78.47±5.02	80.22±2.28	p>0.05
Resp. rate(/min.)	16.01	16.05	P>0.05

The mean pulse rate changes were observed between Butorphanol group and Fentanyl group were statistically nonsignificant ( $p>0.05$ ). The systolic blood pressure changes were statistically nonsignificant in both the groups ( $>0.05$ ). change in mean respiratory rate between two groups

was statistically nonsignificant ( $p>0.05$ ). Diastolic blood pressure changes were statistically nonsignificant in both the groups. ( $p>0.05$ )

#### 4. DISCUSSION:

Low dose bupivacaine although reduces the cardiovascular effects, it was not enough to provide adequate level of sensory blockade and prolonged post-operative analgesia[7&8]. Intrathecal opioids used as adjuncts are capable of producing analgesia of prolonged duration but allow early ambulation of patients because of their sympathetic and motor nerve sparing activities[9].

The principal findings in our study are that intrathecal butorphanol-bupivacaine provides longer duration of sensory blockade and superior analgesia with lesser requirement for rescue analgesia as compared to intrathecal fentanyl bupivacaine.

Local anesthetics such as bupivacaine act mainly by blockade of voltage gated  $\text{Na}^+$  channels in the axonal membrane and presynaptic inhibition of calcium channels[10]. The use of opioids in conjunction with local anesthetic for spinal anesthesia has been associated with decreased pain scores and reduced analgesic requirement in the postoperative period[11&12].

Results of previous studies have demonstrated that intrathecal opioids not only enhance analgesia when added to subtherapeutic (2,7,10) doses of local anesthetics but also prolong recovery.

In our study, there was not much of a difference for the onset of sensory and motor blockade in between the two groups, which is 18,19 correlated with study of Mukherji et. al. and Kumar et al.

The highest sensory level achieved by group-A ( $7.0 \pm 1.8$  min.) and group-B ( $7.2 \pm 2.1$  min.) were comparable (T10).

Two segment regression time was significantly prolonged in group A ( $106 \pm 21$  min.) compared to group B ( $85 \pm 25$  min.) and ( $p<0.001$ ) which is statistically highly significant, which supports the results obtained from Mukherjee A et.al[19] Kumar B et.al.

They concluded that though both Butorphanol and Fentanyl prolonged duration of sensory block of Bupivacaine, Butorphanol is better in terms of longer duration of action. Duration of effective analgesia ( $\text{VAS} \geq 4$ ), was significantly prolonged in group A compared to group B ( $p<0.001$ ). Mean regression time to Bromage 0 motor block was significantly higher in Butorphanol group than Fentanyl group[13,14&15].

In our study we defined hypotension as a decrease of systolic B.P of more than 30% of baseline, 2 patients had hypotension in group A. the patients were treated with one dose of Inj. Ephedrine 3 mg IV each after treating with IV fluids[16&17]. It has been reported that neuraxial administration of opioids with local anesthetics can lead to an 15 increased incidence of hypotension. It has been postulated by Adkinson et al that increased incidence of hypotension following co-administration of fentanyl and lidocaine could be due to higher 16 sensory level achieved. Weight(kg) 48.13 48.06  $>0.05$  Parameters Butorphanol (GP-A) Mean SD Fentanyl (GPB) Mean SD P-Value Highest Sensory level (median range) T7 (T 6-8) T8 (T5 - 8) - Time from injection to highest sensory level (min)  $7.0 \pm 1.8$   $7.2 \pm 2.1$   $P>0.05$  Time of two segment regression (min)  $106 \pm 21$   $85 \pm 25$   $P0.05$  Duration of grade III motor block (min)  $115 \pm 22$   $110 \pm 20$   $P0.05$  Heart rate (/min.)  $78.47 \pm 5.02$   $80.22 \pm 2.28$   $p>0.05$  Resp. rate(/min.) 16.01 16.05  $P>0.05$ [18,19&20].

#### 5. CONCLUSION:

Both Butorphanol and Fentanyl in combination with low dose hyperbaric bupivacaine (15mg) are efficacious in patients undergoing lower limb orthopedic surgeries instead of bupivacaine alone because both opioids having haemodynamic stability with these combinations is good,

effective and prolonged duration of sensory analgesia, Butorphanol with low-dose bupivacaine in spinal anesthesia is better acceptable clinically in terms of characteristics of sensory block, motor block, duration of analgesia and greater hemodynamic stability as compared with Fentanyl-bupivacaine .

## 6. REFERENCES:

- [1] Wang JK, Nauss LA, Thomas JE. Pain relief by intrathecally applied morphine in man. *Anaesthesiology* 1979; 50: 149-151.
- [2] Sukhani R, Stevens RA. Spinal anesthesia. In: Benzon HT, Raja SN, Borsook D, Molloy RE, Strichartz G (eds). *Essentials of Pain Medicine and Regional Anaesthesia*, New York: Churchill Livingstone 1999; pp350-7.
- [3] "WCPI Focus on Pain Series: The Three Faces of Fentanyl". *Aspi.wisc.edu*. Retrieved 2010-07-28.
- [4] Stacey Mayes, PharmD MS, Marcus Ferrone, PharmD BCNSP, 2006. Fentanyl HCl Patient-Controlled Iontophoretic Transdermal System for Pain: Pharmacology The Annals of Pharmacotherapy
- [5] Wang C, Chakrabarti MK, Whitwam JG. Specific enhancement by fentanyl of the effects of intrathecal bupivacaine or nociceptive afferent but not on sympathetic efferent pathway in dogs. *Anesthesiology* 1993; 79: 766-73.
- [6] Chaney MA. Side effects of intrathecal and epidural opioids. *Can J Anaesthesia* 1995; 42:891-903.
- [7] Stoelting RK. Intrathecal morphine: an underused combination for postoperative pain management [editorial]. *Anesth Analg* 1989; 68: 707-9.
- [8] Fournier R, Van Gessel E, Weber A, Gamulin Z. A comparison of intrathecal analgesia with fentanyl or sufentanil in hip replacement. *Anesth Analg* 2000; 90: 918-22.
- [9] Cousins JM, Bridenbough PO. Spinal narcotics and pain relief in acute care. IN: Cousins MJ, Philips JD, eds. *Acute Pain Management* New York:Churchill Livingstone, 1986; pp. 156-7.
- [10] Capogna G, Celleno D, Tagariello V, Loffreda-Maniculli C. Intrathecal buprenorphine for postoperative analgesia in the elderly patient. *Anaesthesia* 1988; 43: 128-30.
- [11] Gear, RW; Miaskowski C; Gordon NC; Paul SM; Heller PH; Levine JD (November 1999). "The kappa opioid nalbuphine produces gender- and dose-dependent analgesia and antianalgesia in patients with postoperative pain". *Pain* 83 (2): 339-45. doi:10.1016/S0304-3959(99)00119-0. PMID 10534607.
- [12] McCrae AF, Wildsmith JA. Prevention and treatment of Hypotension during central neuraxial block. *Br J Anaesth*. 1993; 70:672-80.
- [13] Ben-david B, Frankel R, Arzumonov T, Marchevsky Y, Volpin G. Minidose Bupivacaine-fentanyl Spinal Anaesthesia for surgical repair of Hip fracture of aged. *Anaesthesiology*. 2000; 92: 6-10.
- [14] Lambert DH. Factors influencing spinal anaesthesia. *Int Anesthesiol Clin* 1989; 27: 1320.
- [15] Tawh MO. Mode of action of intraspinal opioids. *Pain rev* 1994; 1: 275-294.
- [16] Terjima K, Onodera H, Koboyashi M, Yamanaka H, Ohno T, Konuma S, et al. efficacy of intrathecal morphine for analgesia following elective cesarean section: comparison with previous delivery. *JNippon Med Sch* 2003; 70: 327-33.
- [17] Butterworth JF 4th, Strichartz GR. Molecular mechanism of LA: A review. *Anaesthesiology* 1990; 72: 71-74.
- [18] Ackerman B, Arwestrom E, post C. Local Anaesthetics potentiate spinal morphine antinociception. *Anaest Anal* 1988; 67: 943-948.

- [19] Mukherjee A, Pal A, Agarwal J, Mehrotra A, Dawar N. Intrathecal nalbuphine as an adjuvant to Subarachnoid block: What is the most effective dose?. *Anesthesia: Essay and Researches* 2011; 5(2)171-175.
- [20] Kumar B, Williams A, Liddell D, Verghese M. Comparison of intrathecal bupivacaine – fentanyl And bupivacaine -butorphanol mixtures for lower limb orthopedic procedures *Anesthesia: Essay and Researches* 2011; 5(2)190-1860.