

Study of clinical profile, risk factors and outcome of congestive cardiac failure patients

Vaibhav Anil Patil¹, Vivek Divakar Chaudhari², Milind Raghunath Waykole³

¹Assistant Professor, Department of Cardiology, Dr Ulhas Patil Medical College & Hospital, Jalgaon, Maharashtra, India.

²Associate Professor, Department of Cardiology, Dr Ulhas Patil Medical College & Hospital, Jalgaon, Maharashtra, India.

³Associate Professor, Department of Cardiology, Dr Ulhas Patil Medical College & Hospital, Jalgaon, Maharashtra, India.

Received Date: 16/01/2023

Acceptance Date: 09/03/2023

Abstract

Background: CCF is a major health problem affecting about 64.3 million persons globally, with an epidemiology that varies widely within and between countries. The international Congestive Heart Failure Study (INTER-CHF) was an international, multicenter cohort study, conducted in sixteen countries in Africa, Asia, the Middle East, and South America with six- and twelve-month follow-ups. **Aim & Objective:** 1.To study risk factors for congestive heart failure. 2.To study outcome of patients of congestive cardiac failure **Methods:** Prospective cross sectional study, **Study setting:** Cardiology Department of tertiary care centre. **Study population:** The study population included all the cases with CCF admitted at a tertiary care center **Sample size:** 100 **Results:** majority of cases were found in above 70 years group e.g.70 followed by 51-60 years age group 22 cases, 61-70,41-50,31-40,21-30 and 18-30 found 17,15,14,5 and 1 case respectively. majority of study participants were Males contributing 60 (60%) and Females 40 (40%). majority of cases had a history of Hypertension 86, followed by Smoking were present in 40 cases, Alcohol history reported by 37 cases, 23 cases were found with DM, Rheumatic heart disease in 17 cases and metabolic syndrome found in 15 cases. Majority of cases complained Dyspnoea on exertion e.g. 89 followed by Orthopnoea in 73 cases, Pedal oedema in 67, Chest pain 56, Palpitation 49, cough 40, syncope 40 and other complained reported by 56 cases. majority of cases discharged 81, DAMA 12 cases and 7 cases death during treatment **Conclusions:** CCF was seen commonly in age group of above 70 years.most common risk factor was HTN. Hospital mortality was 7%

Keywords: CCF, HFpEF, HFrEF, Modifiable risk factors, Outcome

Corresponding Author: Dr. Milind Raghunath Waykole, Associate Professor, Department of Cardiology, Dr Ulhas Patil Medical College & Hospital, Jalgaon, Maharashtra, India.

Email: milindwaykole@gmail.com

Introduction

Congestive cardiac failure is a major health problem affecting about 64.3 million persons globally, with an epidemiology that varies widely within and between countries[1]. The international Congestive Heart Failure Study (INTER-CHF) was an international, multicenter cohort study, conducted in sixteen countries in Africa, Asia, the Middle East, and South America with six- and twelve-month follow-ups [2,3,4].

It was the first major study to systematically acquire data on in- and out-patients with HF in these regions, which were hitherto under-represented in previous global HF studies. It

revealed that mortality rate was highest in Africa (34%), intermediate in southeast Asia (15%), and lowest in China (7%), South America (9%), and the Middle East (9%), and the regional differences persisted after multivariable adjustment [4].

The variations in mortality between the regions were due to an interplay between health-care infrastructure, and environmental and genetic factors [4]. The prevalence is increasing in many countries due to aging societies, increased prevalence of risk factors, and better survival from other cardiovascular diseases.[5-7]) However, the survival rate of HF remains poor, and the health burden from this condition is increasing globally.[8-11]

The prevalence of risk factors such as diabetes, myocardial infarction, and ischemic heart disease has increased in the past few decades, although the survival outcomes from these diseases have also improved.[12,13] Consequently, the prevalence of HF approximately doubled from 0.75% in 2002 to 0.53% in 2013, and the total medical cost increased by about 50% from 2009 to 2013.[14,15] The increase in total medical cost was mostly attributable to the cost of in-hospital care.

Aim and objectives

1. Clinical profile of patients with CCF.
2. To study risk factors for CCF.
3. To study outcome of patients of CCF

Methodology

Study design: Cross Sectional Study.

Study setting: Cardiology tertiary care centre

Study population: The study population included all the cases with Congestive cardiac failure admitted at a tertiary care center

Inclusion Criteria

1. All the classical cases of CCF above the age of 18 years according to Framingham criteria with right sided or left sided independently or in combination (biventricular failure) are included.

Exclusion Criteria

1. Not willing to participate in the study
2. Loss to follow up
3. Patients below the age of 18 years
4. Already diagnosed cases of congenital heart disease

Approval for the study

Written approval from Institutional Ethics committee was obtained beforehand. Written approval of Medicine and Related department was obtained. After obtaining informed verbal consent from all patients with the definitive diagnosis of heart failure admitted to Medicine ward of tertiary care centre such cases were included in the study.

Sample Size: With reference to study by **Mistry BS et al (2019)**¹⁶ He found that the prevalence of IHD among heart failure cases was 64.28%.

Formula for sample size = $4 * P * Q / L^2$

Where P = 64.28%

Q = 100 -64.28 = 35.72

L = Allowable error = 20% (Absolute error)

Sample size = $4 * 64.28 * 35.72 / 163.84=56.05$

Sample size Rounded to = 100

Sampling technique

Convenient sampling technique used for data collection.

All patients admitted in the Cardiology department of tertiary care center with CCF were

included in the study.

Methods of Data Collection and Questionnaire: Predesigned and pretested questionnaire was used to record the necessary information. Questionnaires included general information, such as age, sex, religion, occupation of parents, residential address, and date of admission. Medical history- chief complain, past history, general examination, systemic examination. After taking written and informed consent about enrolment in the study and maintaining adequate privacy and confidentiality, all patients were subjected to a standardized interview. Detailed medical history was taken, and complete general and systemic examinations were done to establish the diagnosis of heart failure and rule out close differentials like renal failure and acute respiratory distress syndrome.

Complete hemogram, urine routine analysis, renal function test (s. urea, s. creatinine), liver function test (s. bilirubin), serum electrolytes (S. Na⁺, S. K⁺) (as an when required), electrocardiogram, chest x-ray, 2d echocardiography, random blood sugar, serum lipid profile, other investigations (as an when required) were done. All the procedures and investigations conducted under direct guidance and supervision of pg guide. Proforma of heart failure notes maintained.

Data entry and analysis

The data were entered in Microsoft Excel and data analysis was done by using SPSS demo version no 21 for windows. The analysis was performed by using percentages in frequency tables and correlation of heart failure. $p < 0.05$ was considered as level of significance using the Chi-square test.

Results And Observations

The present Cross-sectional study was done among 100 CCF cases admitted to tertiary care centre during study period.

Table 1: Distribution of cases according to age (N=100)

Age in years	Frequency	Percentage
18-20	1	1%
21-30	5	5%
31- 40	14	14%
41-50	15	15%
51-60	22	22%
61-70	17	17%
Above 70	26	26%
Total	100	100 (100%)

Above table shows that majority of cases were found in above 70 years group e.g. 70 followed by 51-60 years age group 22 cases, 61-70,41-50,31-40,21-30 and 18-30 found 17,15,14,5 and 1 case respectively.

Table 2: Distribution of cases as per sex (N=100)

Gender	Frequency	Percentage
Male	60	60%
Female	40	40%
Total	100	100 (100%)

Above table shows that majority of study participants were Males contributing 60 (60%) and Females 40 (40%)

Table 3: Distribution of cases according to Risk factors (N=100)

Risk factors	Frequency	Percentage
Hypertension	86	86%
Diabetes mellitus	23	23%
Alcohol	37	37%
Smoking	40	40%
Rheumatic heart Disease	17	17%
Obesity	15	15%

The above table shows majority of cases had a history of Hypertension 86, followed by Smoking were present in 40 cases, Alcohol history reported by 37 cases, 23 cases were found with DM, Rheumatic heart disease in 17 cases and metabolic syndrome found in 15 cases.

Table 4: Distribution according to clinical feature (N=100)

Clinical feature	Frequency	Percentage
Dyspnoea on exertion	89	76%
Orthopnoea	73	73%
Pedal oedema	67	67%
Chest pain	56	56%
Cough	40	40%
Palpitation	49	49%
Syncope	40	40%
Other	56	56%

Majority of cases complained Dyspnoea on exertion e.g. 89 followed by Orthopnoea in 73 cases, Pedal oedema in 67, Chest pain 56, Palpitation 49, cough 40, syncope 40 and other complained reported by 56 cases.

Table 5: Distribution of cases according to outcome (N=100)

Outcome	Frequency	Percentage
Survival/Discharged	81	81%
DAMA	12	12%
Death	7	7%
Total	100	100 (100%)

The above table shows majority of cases discharged 81, DAMA 12 cases and 7 cases death during treatment.

Discussion

The present cross-sectional study was done among 100 cases of CCF admitted to tertiary care centre during study period.

In current study majority of cases were found in above 70 years group e.g. 70 followed by 51-60 years age group 22 cases, 61-70, 41-50, 31-40, 21-30 and 18-30 found 17, 15, 14, 5 and 1 case respectively. Similar result found in Karaye KM et al (2021)¹⁷ He reported that the maximum cases in above 60 years age group. Another study conducted by Harikrishnan S et al (2022)¹⁸ He revealed that the mean age: 59.9 years.

In present study majority of study participants were Males contributing 60 (60%) and Females 40 (40%). similar result observed in the study conducted by Mistry BS et al (2019)¹⁶ He reported that the Males constituted more than half of patients. Another study reported similar result Harikrishnan S et al (2022)¹⁸ He revealed that the 31% women and 69% male.

In current study majority of cases had a history of Hypertension 86, followed by Smoking were present in 40 cases, Alcohol history reported by 37 cases, 23 cases were found with DM, Rheumatic heart disease in 17 cases and metabolic syndrome found in 15 cases. Similar result found in the study by Mistry BS et al (2019)¹⁶ He reported that the Common aetiologies found were ischaemic heart disease (IHD), hypertension (HTN), diabetes mellitus (DM) and valvular heart disease. 50-64 year patients constituted the major age group. IHD and DM were more common in this age group. Rheumatic heart disease (RHD) was seen commonly in 20-34 year of age group.

In current study Majority of cases complained Dyspnoea on exertion e.g. 89 followed by Orthopnoea in 73 cases, Pedal oedema in 67, Chest pain 56, Palpitation 49, cough 40, syncope 40 and other complained reported by 56 cases. Similar result reported by Gupta V et al (2019)¹⁹ He observed that the dyspnoea was most common complain.

In current study majority of cases discharged 81, DAMA 12 cases and 7 cases death during treatment. Similar result found in the study conducted by Mistry BS et al (2019)¹⁶ He reported that the Out of 84 patients, 72 were discharged, 8 took discharge against medical advice and 4 patients expired. In hospital mortality was 5.26%

Conclusions

Heart failure was seen commonly in age group of above 70 years. Majority of cases had a history of Hypertension. Majority of cases complained Dyspnoea. Hospital mortality was 7%.

References

1. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018; 392: 1789–1858. DOI: [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7)
2. Dokainish H, Teo K, Zhu J, et al. Heart failure in low- and middle-income countries: Background, rationale, and design of the INTERnational Congestive Heart Failure Study (INTER-CHF). *Am Heart J*. 2015; 170(4): 627–634. DOI: <https://doi.org/10.1016/j.ahj.2015.07.008>
3. Dokainish H, Teo K, Zhu J, et al. on behalf of the INTER-CHF Investigators. Heart Failure in Africa, Asia, the Middle East and South America: The INTER-CHF study. *Int J Cardiol*. 2016; 204: 133–141. DOI: <https://doi.org/10.1016/j.ijcard.2015.11.183>
4. Dokainish H, Teo K, Zhu J, et al. on behalf of the INTER-CHF Investigators. Global mortality variations in patients with heart failure: Results from the International Congestive Heart Failure (INTER-CHF) prospective cohort study. *Lancet Glob Health*; 2017. DOI: [https://doi.org/10.1016/S2214-109X\(17\)30196-1](https://doi.org/10.1016/S2214-109X(17)30196-1).
5. Heidenreich PA, Trogon JG, Khavjou OA, et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation* 2011;123:933-44.
6. Okura Y, Ramadan MM, Ohno Y, et al. Impending epidemic: future projection of heart failure in Japan to the year 2055. *Circ J* 2008;72:489-91.
7. Curtis LH, Whellan DJ, Hammill BG, et al. Incidence and prevalence of heart failure in elderly persons, 1994-2003. *Arch Intern Med* 2008;168:418-24
8. Sato N, Kajimoto K, Asai K, et al. Acute decompensated heart failure syndromes (ATTEND) registry. A prospective observational multicenter cohort study: rationale, design, and preliminary data. *Am Heart J* 2010;159:949-55.e1.
9. Fonarow GC, Heywood JT, Heidenreich PA, Lopatin M, Yancy CW; ADHERE Scientific Advisory Committee and Investigators. Temporal trends in clinical

- characteristics, treatments, and outcomes for heart failure hospitalizations, 2002 to 2004: findings from Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J* 2007;153:1021-8.
10. Abraham WT, Fonarow GC, Albert NM, et al. Predictors of in-hospital mortality in patients hospitalized for heart failure: insights from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *J Am Coll Cardiol* 2008;52:347-56.
 11. Nieminen MS, Brutsaert D, Dickstein K, et al. EuroHeart Failure Survey II (EHFS II): a survey on hospitalized acute heart failure patients: description of population. *Eur Heart J* 2006;27:2725-36.
 12. Kook HY, Jeong MH, Oh S, et al. Current trend of acute myocardial infarction in Korea (from the Korea Acute Myocardial Infarction Registry from 2006 to 2013). *Am J Cardiol* 2014;114:1817-22.
 13. Ha KH, Kim DJ. Trends in the diabetes epidemic in Korea. *Endocrinol Metab (Seoul)* 2015;30:142-6.
 14. Lee JH, Lim NK, Cho MC, Park HY. Epidemiology of heart failure in Korea: present and future. *Korean Circ J* 2016;46:658-64.
 15. Service KNHI [Internet]. 2014 Sep [cited 2021 October]. Available from: <http://www.nhis.or.kr/bbs7/boards/B0039/3171>
 16. Mistry BS, Pathak K, Trivedi SK. Clinical and etiological profile of heart failure. *Int J Adv Med* 2019;6:439-45.
 17. Karaye KM, Dokainish H, ElSayed A, Mondo C, Damasceno A, Sliwa K, Balasubramanian K, Grinvalds A, Yusuf S; investigators of INTERNATIONAL Congestive Heart Failure (INTER-CHF) Study. Clinical Profiles and Outcomes of Heart Failure in Five African Countries: Results from INTER-CHF Study. *Glob Heart*. 2021 Jul 30;16(1):50. doi: 10.5334/gh.940. PMID: 34381672; PMCID: PMC8323533.
 18. Harikrishnan S, Bahl A, Roy A, Mishra A, Prajapati J, Manjunath CN, Sethi R, Guha S, Satheesh S, Dhaliwal RS, Sarma M, Ganapathy S, Jeemon P; NHFR investigators. Clinical profile and 90 day outcomes of 10 851 heart failure patients across India: National Heart Failure Registry. *ESC Heart Fail*. 2022 Aug 10. doi: 10.1002/ehf2.14096. Epub ahead of print. PMID: 36214477.
 19. Gupta V, Redkar N, Jena A. A Study of Clinical Profile and Outcome of Acute Heart Failure in Elderly Patients. *J Assoc Physicians India*. 2019 Nov;67(11):55-58. PMID: 31793270.