

# Repeated Pneumonia in Children with Congenital Heart Disease in Tertiary care Hospital at Jharkhand

<sup>1</sup>Dr Partha Kumar Chaudhuri, Dr Divya Singh <sup>2</sup>, Dr Bhuwan Kumar Singh <sup>3</sup>

<sup>1</sup> MD, Associate Professor, Department of Paediatrics, Rajendra Institute of Medical Sciences, Ranchi.

<sup>2</sup> MD, Medical Officer, Department Of Paediatrics, Rajendra Institute of Medical Sciences, Ranchi.

<sup>3</sup> MD, Senior Resident, Department of Paediatrics, Rajendra Institute of Medical Sciences, Ranchi.

**Corresponding Author:**  
**Partha Kumar Chaudhuri**  
pchaudhuri73@yahoo.com

## Abstract

**Background:** Functions of the respiratory and cardiovascular systems have a close relationship. Congenital heart disease (CHD) affects the respiratory system and result in respiratory morbidities. Repeated pneumonia is one such morbidity. **Aim:** To assess the incidence of CHD in patients presenting with repeated pneumonia and the incidence of different types of CHD in patients detected with congenital heart defects. **Materials and Methods:** A total of 100 patients with repeated pneumonia were studied (N = 100), which included 60 males and 40 females. All the patients were subjected to clinical examination and were made to undergo chest X-ray, electrocardiography, and 2D echocardiography with colour Doppler to detect CHD. **Results:** Of the 100 patients, 43 patients (23 [53.4%] males and 20 [46.5%] females) with repeated pneumonia were found to have CHD. **Conclusions:** The results reveal that CHD is a major cause for recurrent LRTI. Considering this, it would be prudent to screen all children presenting with repeated pneumonia for CHD.

**Keywords:** Congenital heart disease, Children, Pneumonia.

## I. Introduction

Functions of the respiratory and cardiovascular systems have a close relationship. Congenital heart disease (CHD) affects the respiratory system and result in respiratory morbidities. Recurrent respiratory infection is one such morbidity. In the presence of congenital anomalies of the circulatory system, the ability of the heart to increase systemic and/or pulmonary blood flow is often limited; arterial partial pressure of oxygen may be decreased by shunt lesions, affecting oxygen delivery to the tissues. Often the circulatory derangement also places stress on the respiratory system itself, resulting in signs and symptoms that mimic primary respiratory disease. Recurrent lower RTI (LRTI) refers to 2 or more hospitalization in 6 months or 3 hospitalizations for RTI in any time frame .1-3 The aetiology of RTI is numerous, infections (viral, bacterial) being the most common. However, underlying CHD may be the predisposing factor for recurrent RTI. Although, congenital defects in the small left-to-right shunts do have symptoms, such as recurrent LRTI and failure to thrive, during early childhood, screening this subset of patients for CHD is worthwhile. Early detection and appropriate management of CHDs can provide a child ample time for catch-up growth, reduce the morbidity and mortality risk associated with each episode of LRTI, significantly reduce the financial burden on the family, and prevent long-term morbidities. Early and accurate diagnosis of CHD in children presenting under the disguise of recurrent LRTI requires prompt, effective, and systematic approaches including detailed medical history collection and thorough clinical examination.

## Aim

To assess the incidence of CHD in patients presenting with recurrent LRTI and the incidence of different types of CHD in patients detected with congenital heart defects.

## Materials And Methods

This study was conducted at the Department of Paediatrics, RAJENDRA INSTITUTE OF MEDICAL SCIENCES (RANCHI, JHARKHAND, INDIA) between September 2017 and August 2018 .

**Inclusion Criteria**

The patients were initially said to have recurrent LRTI based on the following Indian Academy of Paediatrics (IAP) criteria.

1. At least 2 episodes of pneumonia occurring in 1 year or 3 episodes of pneumonia occurring over any period of time
2. Between 2 different episodes of recurrent pneumonia the individual recovers completely but without radiologic improvement

**Exclusion Criteria**

1. Those unwilling to undergo chest X-ray, electrocardiography (ECG), and echocardiography (ECHO)
2. Those already operated for CHD
3. Known cases of bronchial asthma

**Study Procedure** Detailed medical history of all the 100 patients was collected. All patients underwent thorough clinical examination followed by investigative workup. Chest X-ray, ECG, and 2D ECHO with color Doppler were performed by a paediatrician and then the finding was confirmed by a cardiologist. Routine investigations such as haemoglobin (Hb) level, total count (TC), different count (DC), erythrocyte sedimentation rate (ESR) was also conducted.

**Results**

A total of 100 patients with recurrent LRTI were studied (N = 100), which included 60 males and 40 females. Of the 100 patients, 43 patients (23 [53.4%] males and 20[46.5%] females) with recurrent LRTI were found to have CHD

**Table 1.** Type of CHD in patients with repeated pneumonia

Type Of Child	No. Of Patients	Percentage
Acyanotic	39	90.7
Cyanotic	4	9.3
Total	43	100

**Table 2.** Types of ACHD in patients with repeated pneumonia

Type of children	No. Of Patients	Percentage
VSD	17	43.6
ASD	11	28.2
PDA	7	17.9
VSD+ASD	3	7.7
MVP	1	2.5
Total	39	100

**Table 3.** Types of CCHD in Patients with repeated pneumonia

Types of children	No. Of Patients	Percentage
TOF	2	66.66
Single Ventricle	1	33.33
Total	3	100

**Table 4.** Age distribution of patients with recurrent LRTI

AGE	Percentage
<1 y	27
1-5 y	45
6-10 y	21
>10 y	7

**Table 5.** Age distribution of patients with CHD

AGE	Percentage
<1 y	35
1-5 y	40
6-10 y	15
>10 y	10

**Discussion**

CHD is one of the underlying causes for repeated pneumonia, which when detected early and treated appropriately can prevent the child from developing irreversible and untreatable conditions such as pulmonary hypertension, Eisenmenger syndrome, thromboembolic phenomenon, and sudden death. In this study, the incidence of CHD among children was found to be 43%. This is significantly higher than its incidence among the general population (6-8%). This signifies that CHD is a major cause for recurrent LRTIs. Hence, it is important to identify children with recurrent LRTI and screen them for any underlying CHD using through physical and clinical examinations. Chandramouli<sup>4</sup>, in his study on 2613 children < 24 months old with CHD, reported that bronchiolitis (LRTI) was the commonest cause of hospitalization (54.1%). Gupte et al<sup>5</sup> reported that CHDs are present in 36% patients with recurrent LRTI.

***Type of CHD in patients with recurrent LRTI***

Majority of children had acyanotic congenital heart disease (ACHD) (39/43). Similar results were reported in studies conducted by Shreshta et al,<sup>6</sup> and Suresh et al,<sup>7</sup> in India. Three cases were found to have CCHD, of which 2 cases had Tetralogy Of Fallot (TOF). One child has single ventricle.

***Age distribution***

Most of the children with recurrent LRTI were aged between 1 and 5 years and most children with CHD were also in age group of 1 to 5 years. Thus the incidence of CHD is common in that age group with statistically significant ( $P < 0.03$ ).

**Conclusion**

The result show that the CHD is major cause for recurrent LRTIs. Considering this, it would be necessary to screen all children presenting with repeated pneumonia for CHD in clinical practice

**References**

- [1]. Behrman RE, Kliegman RM, Jenson HB, eds. congenital heart disease. In: Nelson Textbook of Pediatrics. 19th ed. Harcourt Aisa Pvt Ltd; 2000:1549-1605.
- [2]. Park KM. Paediatric Cardiology for Practitioners. 5th ed. Elsevier Health Sciences; 2007:119-326.
- [3]. Subramanyam L. Recurrent respiratory tract infection – an approach. Indian J PractPediatr. 2012;14(3):245-257.
- [4]. Chandramouli B. The neonate with congenital heart disease: diagnosis and management. Indian J Pediatr 1991;58(4):453-469.
- [5]. Gupta S , Saini G. Congenital heart disease: Clinicoechocardiographic profile in children. Asian J PediatrPract. 2004;8(2):30-34.
- [6]. Shreshta NK, Padmavati S. Congenital heart disease in Delhi school children. Indian J Med Res. 1980;72:403-407
- [7]. Suresh V, Rao AS, Yavagal ST. Frequency of various congenital heart disease: analysis of 3790 consecutively catheterized patients. Indian Heart J. 1995;47(2): 125-128