

Original research article**An analytical cross-sectional assessment of the association of sensorineural hearing loss in type 2 diabetes mellitus patients****¹Dr. Hari Prasad Garudasu, ²Umavani Hiranmayee Kondabattini**¹Assistant Professor, Department of ENT, Mamata Medical College, Khammam, Telangana, India²Assistant Professor, Department of General Medicine, Mamata Medical College, Khammam, Telangana, India**Corresponding Author:**Dr. Hari Prasad Garudasu (harigarudasu@gmail.com)**Abstract**

Aim: The aim of the present study was to find the prevalence of SNHL in type 2 diabetes patients and to find the effect of duration and control of diabetes on hearing loss.

Methods: The present analytical cross-sectional study was conducted in Department of Otorhinolaryngology, Mamata Medical College, Khammam, for period of 1.5 years after obtaining the ethical clearance from the Institutional Ethical Committee. The present study was conducted on 200 type 2 diabetes patients and age and gender matched 200 non-diabetic controls in the age group of <50 years.

Results: In the present study, 68% were females and 32% were male. The mean age of patients was 44.16, mean fasting blood sugar was 130.60±28.62, mean PP blood sugar was 172.18±42.28 and mean HbA1c was 7.40±0.45. Mean age of patients was 36.60, mean fasting blood sugar was 108.02±3.16, mean post-prandial blood sugar was 126.54±9.99 and mean HbA1c was 6.004±0.220. It showed that only 50 patients had normal hearing and 150 patients had SNHL whereas 32 patients in this group had minimal hearing, 38 had mild SNHL, 48 had moderate, 12 had severe SNHL and 20 patients had profound SNHL. 140 patients had DPOAE refer whereas 60 patients had DPOAE pass.

Conclusion: This study concluded that the diabetics are at definite risk of developing auditory dysfunction, therefore it is recommended that all newly diagnosed diabetic patients should undergo a complete audiological evaluation at the time of diagnosis and a regular half yearly or yearly follow up is warranted for early detection of damage to auditory functions. Although factors other than diabetes contribute to hearing loss, early glycemic control for type 2 diabetic patients may reduce the incidence rate of this disease.

Keywords: Sensorineural hearing loss, diabetes mellitus

Introduction

There is little awareness of hearing loss as a possible comorbid condition associated with type 2 diabetes among persons with diabetes as well as healthcare professionals, despite several studies having demonstrated the link between the two clinical conditions [1]. More than 43% of diabetes patients are likely to have some degree of hearing impairment related or unrelated to chronic hyperglycemia [2]. There is a requirement for further studies for the exploration of the relationship between diabetes and hearing abnormalities. The presence of hearing defects among diabetes patients could be related to hyperglycemia or other associated conditions like decreased immunity that may predispose one to ear infections involving the external, middle, or internal ear [3].

Diabetic neuropathy, which affects the eighth cranial nerve directly or at the cochlear level, may present with variable degrees of hearing loss, while chronic infections like malignant otitis externa that affect the external ear are usually observed among diabetes patients and usually are associated with structural damage [3]. Sufficient data support that hearing loss is one of the commonly occurring diseases in diabetes patients that could affect quality of life and lead to hearing disabilities and psychological depression [2]. Some studies have found positive correlations between hearing loss and diabetes duration or the degree of metabolic control. Although the findings of studies focusing on the relationship between hearing acuity and the degree of hyperglycemia by diabetes type are conflicting, [4] several risk factors have been identified as being associated with hearing loss in diabetes patients [5].

In addition to the cardiovascular outcomes and the long term damage to the kidneys, eyes, and nervous system caused by diabetes [6], this disease was also associated with alterations in hearing function in a meta-analysis of 18 clinical and epidemiological studies. [7] The pathological support to this association may be related to an increase in capillary lesions in the cochlea, more specifically in the stria vascularis

and basilar membrane. Other studies have also reported a reduction in the number of spiral ganglion neurons. These differences could be related to the duration of diabetes and to comorbidities that could affect the inner ear [8]. Although, several studies have investigated the effects of diabetes on hearing loss (HL), the relationship between these two conditions remains unclear. Some studies have suggested that diabetes may cause sensorineural hearing loss (SNHL) [7-10], whereas others have failed to find an association [11-15].

The aim of the present study was to find the prevalence of SNHL in type 2 diabetes patients and to find the effect of duration and control of diabetes on hearing loss.

Materials and Methods

The present analytical cross-sectional study was conducted in Department of Otorhinolaryngology, Mamata Medical College, Khammam for period of 1.5 years after obtaining the ethical clearance from the Institutional Ethical Committee. The present study was conducted on 200 type 2 diabetes patients and age and gender matched 200 non-diabetic controls in the age group of <50 years. Study group included patients who were diagnosed as type 2 DM attending diabetic clinic in Endocrinology OPD while the control group comprised of age and sex matched non-DM individuals. Both groups were subjected to thorough audiological evaluation.

Inclusion criteria

Inclusion criteria for current study were; biochemically proven hyperglycemic patients, age less than or equal to 50 years of both genders.

Exclusion criteria

Exclusion criteria for current study were; patients with any other systemic illness or metabolic disorder, gestational, patients on ototoxic drugs or any ear surgery done/occupational exposure to noise, noise induced hearing loss, patients with CSOM and CHL and patients with congenital SNHL.

Methodology

The present study was a comparative study with population of 100 diagnosed patients of diabetes mellitus and control group consists of 100 non-DM patients. All these individuals were evaluated with a detailed clinical examination and appropriate investigation. All these individuals were evaluated with a detailed clinical examination and appropriate investigation, such as PTA, BERA and DPOAE.

Statistical analysis

Data was entered in Microsoft Excel Spreadsheet. Continuous variables were summarised as mean and standard deviation (SD). Categorical variables were summarised as frequency and percentage. The reported p values were based on analysis, in which $p < 0.5$ was considered significant.

Results

Table 1: Demographic details

| Gender | N | % |
|--------------------------------|---------------------------------|--------------------------------|
| Male | 64 | 32 |
| Female | 136 | 68 |
| Total | 200 | 100.0 |
| Parameters | Diabetic group | Non-diabetic group |
| Mean age (years) | 44.16 ±6.30 | 35.60±8.60 |
| Mean fasting blood sugar | 130.60±28.62 (Range 81-253) | 108.02±3.16 (Range 100-118) |
| Mean post prandial blood sugar | 172.18±42.28 (Range 107-346) | 126.54±9.99 (Range 108-148) |
| Mean HbA1c | 7.40±0.45 (Range 5.4-9.6) | 6.004±0.220 (Range 5.3-6.5) |

In the present study, 68% were females and 32% were male. The mean age of patients was 44.16, mean fasting blood sugar was 130.60±28.62, mean PP blood sugar was 172.18±42.28 and mean HbA1c was 7.40±0.45. Mean age of patients was 36.60, mean fasting blood sugar was 108.02±3.16, mean post-prandial blood sugar was 126.54±9.99 and mean HbA1c was 6.004±0.220.

Table 2: Grade of hearing loss in study participants

| PTA | N | % |
|-------------|----|----|
| Normal | 50 | 25 |
| MinimalSNHL | 32 | 16 |

| | | |
|--------------|-----|-------|
| MildSNHL | 38 | 19 |
| ModerateSNHL | 48 | 24 |
| SevereSNHL | 12 | 6 |
| ProfoundSNHL | 20 | 10 |
| Total | 200 | 100.0 |

The result showed that only 50 patients had normal hearing and 150 patients had SNHL whereas 32 patients in this group had minimal hearing, 38 had mild SNHL, 48 had moderate, 12 had severe SNHL and 20 patients had profound SNHL.

Table 3: Distribution of study patients according to DPOAE

| DPOAE | N | % |
|-------|-----|-------|
| Refer | 140 | 70 |
| Pass | 60 | 30 |
| Total | 200 | 100.0 |

140 patients had DPOAE refer whereas 60 patients had DPOAE pass.

Table 4: Comparative BERA results of diabetic and non-diabetic groups

| Wavelatencies | Intensity (dBnHL) | Non-diabetic group Mean \pm SD | Diabetic group Mean \pm SD (ms) | P value |
|---------------|-------------------|----------------------------------|-----------------------------------|---------|
| V | 70 | 6.04 \pm 0.20 | 6.36 \pm 1.50 | <0.001 |
| I-V | 70 | 3.20 \pm 0.04 | 3.70 \pm 0.24 | <0.001 |
| V | 80 | 5.55 \pm 0.25 | 6.50 \pm 1.36 | 0.003 |
| I-V | 80 | 4.16 \pm 0.12 | 4.48 \pm 0.40 | 0.030 |
| V | 90 | 5.35 \pm 0.20 | 6.46 \pm 1.48 | <0.001 |
| I-V | 90 | 4.76 \pm 0.21 | 5.55 \pm 0.60 | <0.001 |

Wave V grossly delayed in DM group as compared to non-DM with all frequencies (i.e. 70, 80 and 90) whereas interpeak latencies of wave I-V at 70 db frequency is almost equal in DM and non-DM group but it increase with increase in frequencies (i.e. 80 and 90 db).

Discussion

Type 2 diabetes mellitus (DM) is a syndrome of chronic hyperglycemia due to relative deficiency of insulin, resistance to insulin or both [16]. Type 2 diabetes mellitus occur as a result of obesity and lack of exercise [17]. Some people are genetically more prone to diabetes [18]. Globally as of 2015, it was estimated that there were 392 million people with type 2 diabetes making up about 90% of diabetes cases [19, 20]. This is equivalent to about 6% of the world's population [20]. The aim of the present study was to find the prevalence of SNHL in type 2 diabetes patients and to find the effect of duration and control of diabetes on hearing loss.

In the present study, 68% were females and 32% were male. The mean age of patients was 44.16. Study by Ashish *et al* conducted on diabetic patients consisting of similar age group between 18-50 years of age [21]. It was found in their study that majority of them had minimal hearing loss followed by mild hearing loss. 30% of patients had normal hearing in both ears. Rajendran *et al* conducted a similar study in the age group 40-50 years and found that number of people affected with SNHL among the diabetes is 73.3% when compared to that of controls [22].

The mean fasting blood sugar was 130.60 \pm 28.62, mean PP blood sugar was 172.18 \pm 42.28 and mean HbA1c was 7.40 \pm 0.45. Mean age of patients was 36.60, mean fasting blood sugar was 108.02 \pm 3.16, mean post-prandial blood sugar was 126.54 \pm 9.99 and mean HbA1c was 6.004 \pm 0.220. The result showed that only 50 patients had normal hearing and 150 patients had SNHL whereas 32 patients in this group had minimal hearing, 38 had mild SNHL, 48 had moderate, 12 had severe SNHL and 20 patients had profound SNHL. This study corresponds to the study conducted by Ashish *et al* who had similar results as of our study. In his study he found only 30% of patients had normal hearing and 70% had SNHL. [21-23] Sunkun *et al* conducted a similar study on DM patients and found in his study that 82% had SNHL. 18 Rajendran S, *et al*, [21] conducted a similar study on DM patients and in his results 73.3% of DM patients had SNHL as compared to 6.7% of that of non-DM patients. Dadhich conducted a similar study and it was found in this study that 73% patients had SNHL [24]. Tiwari conducted a similar study and he also found 76.8% patients of DM had SNHL [25].

Wave V grossly delayed in DM group as compared to non-DM with all frequencies (i.e. 70, 80 and 90) whereas interpeak latencies of wave I-V at 70 db frequency is almost equal in DM and non-DM group but it increase with increase in frequencies (i.e. 80 and 90 db). It was found in present study that that latency of wave V was grossly delayed at all frequencies whereas interpeak latencies of wave I-V was almost normal at 70 db frequency but it increase with increase in frequencies (i.e. at 80 and 90 db). This study corresponds to study done by Joshi *et al* who found that absolute latencies of BERA were significantly delayed in for waves II and V and significantly delay was notice in interpeak latencies (IPL) of wave I to III and wave I to V [26]. Chhaya *et al* found that there was significant difference for BERA

abnormalities (for 70, 80 and 90 db) in study group when compared to controls [27]. Siddarth *et al* done a study on topic, brainstem auditory response in type 2 DM [28].

Conclusion

This study concluded that the diabetics are at definite risk of developing auditory dysfunction, therefore it is recommended that all newly diagnosed diabetic patients should undergo a complete audiological evaluation at the time of diagnosis and a regular half yearly or yearly follow up is warranted for early detection of damage to auditory functions. Although factors other than diabetes contribute to hearing loss, early glycemic control for type 2 diabetic patients may reduce the incidence rate of this disease. Other comorbidities including hypertension and hyperlipidemia must be considered in the formulation of strategies to reduce the risk of hearing loss. Awareness must be raised on the significance of hearing loss as a commonly occurring comorbidity in diabetes among healthcare providers. Hearing acuity screening should be a part of routine screening for diabetes patients, and should be conducted on a regular basis to avoid the devastating consequences of this often-overlooked medical condition.

References

1. Hlayisi VG, Petersen L, Ramma L. High prevalence of disabling hearing loss in young to middle-aged adults with diabetes. *International Journal of Diabetes in Developing Countries*. 2019 Jan 1;39:148-53.
2. Pemmaiah KD, Srinivas DR. Hearing loss in diabetes mellitus. *International Journal of Collaborative Research on Internal Medicine & Public Health*. 2011;3(10).
3. Gazzaz ZJ, Makhdom MN, Dhafar KO, Maimini O, Farooq MU, Rasheed A. Patterns of otorhinolaryngological disorders in subjects with diabetes. *IIUM Medical Journal Malaysia*. 2011 Dec 1;10(2).
4. Panchu P. Auditory acuity in type 2 diabetes mellitus. *International journal of diabetes in developing countries*. 2008 Oct;28(4):114.
5. Gutierrez J, Jimeno C, Labra PJ, Grullo PE, Cruz TL. Prevalence of Sensorineural Hearing Loss and its Association with Glycemic Control in Filipino Patients with Diabetes at the Philippine General Hospital. *Journal of the ASEAN Federation of Endocrine Societies*. 2016 Nov 6;31(2):137-.
6. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2004;27(suppl 1):S5-10.
7. Akinpelu OV, Mujica-Mota M, Daniel SJ. Is type 2 diabetes mellitus associated with alterations in hearing? A systematic review and meta-analysis. *Laryngoscope*. 2014;124(3):767-76.
8. Akinpelu OV, Ibrahim F, Waissbluth S, Daniel SJ. Histopathologic changes in the cochlea associated with diabetes mellitus--a review. *Otol Neurotol*. 2014;35(5):764-74.
9. Mitchell P, Gopinath B, McMahan CM, Rochtchina E, Wang JJ, Boyages SC, Leeder SR. Relationship of Type 2 diabetes to the prevalence, incidence and progression of age-related hearing loss. *Diabetic Medicine*. 2009 May;26(5):483-8.
10. Horikawa C, Kodama S, Tanaka S, Fujihara K, Hirasawa R, Yachi Y, *et al*. Diabetes and risk of hearing impairment in adults: a meta-analysis. *The Journal of Clinical Endocrinology & Metabolism*. 2013 Jan 1;98(1):51-8.
11. Harner SG. Hearing in adult-onset diabetes mellitus. *Otolaryngol Head Neck Surg*. 1981;89(2):322-7.
12. Miller JJ, Beck L, Davis A, Jones DE, Thomas AB. Hearing loss in patients with diabetic retinopathy. *Am J Otolaryngol*. 1983;4(5):342-6.
13. Hodgson MJ, Talbott E, Helmkamp JC, Kuller LH. Diabetes, noise exposure, and hearing loss. *J Occup Med*. 1987;29(7):576-9.
14. de España R, Biurrun O, Lorente J, Traserra J. Hearing and diabetes. *ORL J Otorhinolaryngol Relat Spec*. 1995;57(6):325-7.
15. Dalton DS, Cruickshanks KJ, Klein R, Klein BE, Wiley TL. Association of NIDDM and hearing loss. *Diabetes Care*. 1998;21(9):1540-4.
16. Anjana RM, Ali MK, Pradeepa R, Deepa M, Datta M, Unnikrishnan R, Rema M, Mohan V. The need for obtaining accurate nationwide estimates of diabetes prevalence in India--rationale for a national study on diabetes. *The Indian journal of medical research*. 2011 Apr;133(4):369.
17. Diabetes blue circle symbol. *International Diabetes Federation*.
18. Causes of diabetes. *National Institute of Diabetes and Digestive and Kidney Diseases*.
19. Melmed S, Polonsky KS, Larsen P. *Williams textbook of endocrinology*. 12th ed. Philadelphia: Elsevier/Saunders; c2005:1371-35.
20. Vos T, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, Abera SF, Aboyans V. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*. 2017 Sep 16;390(10100):1211-59.
21. Rajendran S, Anandhalakshmi, Mythili B, Viswanatha R. Evaluation of the incidence of sensorineural hearing loss in patients with type 2 diabetes mellitus. *Int J Biol Med Res*.

- 2011;2(4):982-7.
22. Lerman-Garber I, Cuevas-Ramos D, Valdés S, Enríquez L, Lobato M, Osornio M. Sensorineural hearing loss a common finding in early-onset type 2 diabetes mellitus. *EndocrPract.* 2012;18(4):549-57.
 23. Sunkum AJK, Pingile S. A clinical study of audiological profile in diabetes mellitus patients. *Eur Arch Otorhinolaryngol.* 2013;270(3):875-9.
 24. Dadhich S, Jha SG, Sinha V, Samanth TU. A prospective, observational study of incidence of sensory neural hearing loss in diabetes mellitus patients. *Indian J Otol.* 2018;24(2):80-2.
 25. Tiwari A, Mudhol RS. Prevalence of sensorineural hearing loss among type-II diabetes mellitus patients attending KLES Dr. Prabhakar Kore Hospital and MRC: A cross-sectional study. *Indian J Health Sci Biomed Res.* 2018;11:165-9.
 26. Joshi KD, Galagali JR, Kanzhuly MK, Singh ID. A comparative study on effects of diabetes on auditory functions as measured by BERA and DPOAE. *Int J Otorhinolaryngol Head Neck Surg.* 2017;3:347-53.
 27. Batham C, Choudhary AK, Yousuf PS. Brainstem auditory evoked responses with duration of type-II diabetes mellitus. *Ann Med Health Sci Res.* 2017;7:40-5.
 28. Suresh S, Ramlan S, Somayaji G, Sequeira N. Brainstem auditory responses in type-2 diabetes mellitus. *Int J Oto Head Neck Surg.* 2018;4(2):45-9.