

Original Article

“To Study Thalassemia Major Patients And Their Hematological Profile In Tertiary Care Center”

Dr. Chandan Kachhwaha¹, Dr. Meghna Nema², Dr. Amrita Chauhan³,
Dr. Amit Kumar Singh^{4*},

¹Ex. Senior Resident, Department of Pediatrics, MGM Medical College Indore.

²DM Neonatology Senior Resident, Department of Neonatology, Govt Medical College Thiruvananthapuram, Kerala

³Ex. Senior Resident Department of Pediatrics MGM Medical College Indore.

^{4*}Ex. Senior resident, Department of Pediatrics, All India institute Of Medical Sciences Bhopal (M.P.), dramitsingh58@gmail.com

***Corresponding Author:** Dr. Amit Kumar Singh

*Ex. Senior resident, Department of Pediatrics, All India institute Of Medical Sciences Bhopal (M.P.), dramitsingh58@gmail.com

Received 25 July

Revised 28 July

Published 30 July

Abstract-

Introduction- Thalassemia is an inherited disorder of autosomal recessive gene caused by impaired synthesis of one or more globin chains. The impairment alters production of normal hemoglobin (Hb). Thalassemia causes varying degrees of anemia, which can range from significant to life threatening. People of Mediterranean, Middle Eastern, African, and Southeast Asian descent are at higher risk of carrying the genes for thalassemia. Thalassemia causes varying degrees of anemia, which can range from significant to life threatening. India is a large Southeast Asian country with a population of over one Billion. An estimated 1-3% of the population are carriers of beta thalassemia, a figure rising up to 17% in some ethnic groups.

Aims and objectives- This is a prospective observational cohort study done in pediatric wards of two hospitals C. R. Gardi Hospital (C.R.G.H.) and associated hospitals, from September 2016 may 2018. Main aim of this study is- **“to study Thalassemia major patients and their hematological profile in tertiary care center”**

Results- Out of 60 patients of thalassemia were enrolled and their demographic hematological profile was taken. The mean age was 10.2 yrs, 75% were males and 81% were Hindus ,40% belonged to upper lower class .

Most common clinical features were (97%) icterus followed by (90%)pallor,80% of thalassemia patients had moderate to severe hepatomegaly. Mean Hb was 8.5±0.9gm/dl, 67% had severe anemia, Mean MCV level was 73 ±2 μm³,Mean MCH and MCHC level were 20±2pg/cell,28±2gHb/dl.

Mean ferritin level was 1281.8 ± 219.9 ng/dl , 45% of thalassemic patients had serum ferritin level between 1000 to 2000 ng/dl followed by 39% of patients had serum ferritin level more than 2500 ng/dl. 86% had euglycemia and 10% had hyperglycemia when blood sugar was tested randomly.

90% of thalassemic patients had normal glucose tolerance test followed by 6.5% of thalassemic patients had impaired glucose tolerance test, while 3.3 % patients came under diabetic range.

81% of thalassemic patients had normal serum calcium level while 18% had hypocalcemia, 93% of thalassemic patients had normal serum T4, and TSH level ,1.6% of patients had subclinical hypothyroidism and only 4% of thalassemic patients had overt hypothyroidism.

Keywords- ferritin, glucose tolerance test, hypocalcemia.

INTRODUCTION

Thalassemia is caused by mutations that decrease hemoglobin synthesis and red cell survival. Thalassemia is caused by decreased or absent production of one type of globin chain; either alpha or beta globin chain. These hematologic disorders range from asymptomatic to severe anemia that can cause significant morbidity and mortality.

It was first recognized clinically in 1925 by Dr. Cooley, who described a syndrome of anemia with microcytic erythrocytes. Then it was called Cooley's anemia. Later Wipple and Bradford renamed this disease as "Thalassemia". Because it was found in the region of the Mediterranean Sea (thalasa is an old Greek word for sea), (Cooley T B, 1946).

Beta-thalassemia includes three main forms: Thalassemia major variably referred to as "Cooley's Anemia" and "Mediterranean Anemia", Thalassemia intermedia and Thalassemia minor also called "beta-thalassemia carrier", "beta-thalassemia trait" or "heterozygous beta-thalassemia".

Beta thalassemia is a disorder that reduces the production of hemoglobin. Hemoglobin is the iron-containing protein in red blood cells that carries oxygen to cells throughout the body.

In people with beta thalassemia, low levels of hemoglobin lead to a lack of oxygen in many parts of the body. Affected individuals also have decreased red blood cells (anemia), which will manifest as pale skin, weakness, fatigue and serious complications.

It is estimated that 1.5% of the world's population are carriers of β - thalassemia with an estimated 60,000 new carriers born each year (Rathod DA, 2007).⁽⁴³⁾ Southeast Asia accounts for about 50% of the world's carriers while Europe and the Americas jointly account for 10-13% of the world carriers (Angastinotis M, 2005).⁽³⁷⁾

India is a large Southeast Asian country with a population of over one Billion. An estimated 1-3% of the population are carriers of beta thalassemia, a figure rising up to 17% in some ethnic groups (Sukumaran and Master, 1973; Modell and Petrou, 1983).^(7,12)

In India, prevalence of Thalassemia is very high in Punjabis, Sindhs, Gujaratis, Bengalis, Parsee, Lohana and certain tribal communities, and in Northern, Western and Eastern part, while it is much less in the south of India, (Shah et al. 2004).⁽³⁴⁾

About 6,000 children are born with thalassemia major each year, more than 30% of birth with a major thalassemia syndrome in South East Asia (Modell and Petrou, 1983).⁽¹²⁾

Overtime, an influx of iron-containing hemoglobin from chronic blood transfusions can lead to a buildup of iron in the body, resulting in liver, heart, and hormone problems,(Gene Reviews 2015.⁽⁶⁵⁾ The aims and objective of this study is to study the clinical features , complications in Beta-thalassemia patient and outcome from Ujjain district and so that timely intervention can be done to prevent complications.

AIMS AND OBJECTIVES

This is a prospective observational cohort study done in pediatric wards of two hospitals C. R. Gardi Hospital (C.R.G.H.) and associated hospitals, from September 2016 may 2018. Main aim of this study is- **“to study Thalassemia major patients and their hematological profile in tertiary care center”**

Study participants

Inclusion criteria:

- 1) Children with diagnosis of thalassemia
- 2) Age group 6 months and above
- 3) Thalassemia children who are on regular blood transfusion with or without iron chelation therapy

Exclusion criteria: Children with other causes of anemia like nutritional anemia, aplastic anemia, sickle cell anemia.

Sample size

To calculate sample size it was assumed that 50% of patients of Thalassemia will have at least one known complications thus to detect at least 50% of complication with a difference of +/- 20% around the assumed complications rate of 50% with a power of 80 two sided alpha 0.05 estimated minimum sample size is 103.

Data collection: Children who fulfill the inclusion criteria for the study will be selected. After proper clinical examination and preliminary investigation, the child will be subjected to clinical outcome and complication. The data will collect on proper proforma.

The data thus collected will be subjected to statistical analysis.

Pre and post transfusion hemoglobin and serum ferritin level , complete hemogram, Hb electrophoresis, peripheral blood smear ,urine urobilinogen are done as a part of standard care for thalassemia patients.

Serum ferritin level will be estimated by chemiluminescent immunometric assay⁷

Haemoglobin estimation will be done by Beckmen Coulter machine using cyanmethemoglobin automated method.

STATISTICAL ANALYSIS

The data was entered in Epidata Entry (version 3.1) and then transferred to Stata 10.0 (Stata Corp. College Station, Texas, USA) software for statistical analysis. Frequency and percentages are presented for categorical data. Chi square test was used for measuring association between different categories. Binary logistic regression were applied and calculate odd ratio. A *P* value less than or equal to 0.05 was considered significant. t-test was used for comparing the mean duration of stay.

OBSERVATION AND RESULTS

- 50% of thalassemic patients belonged to 5-10yrs of age,28% of patients belonged to 6months-5 yrs and 22% belonged to >10yrs of age.
- 75% of thalassemic patients were males.
- 81% of thalassemic patients belonged to hindu religion and 19% belonged to population.

- 40% of thalassemic patients belonged to upper lower class, 38% patients belonged to lower class, 12% of patients belonged to lower middle class and only 10% patients belonged to upper middle class.
- This table shows various clinical features with which patients presented icterus (97%), pallor (90%), facial changes (79%), loss-of-appetite (60%), lethargy (60%), skin-pigmentation (58%), breathlessness (33%), skeletal changes (30%), diarrhea (27%), fever (14%), oliguria (12%) respiratory distress (17%), hematemesis (16%), dark colored urine (11%), vomiting (7%), edema (2%).
- 50% of thalassemia patients were presented with severe hepatosplenomegaly followed by 30% patients with moderate hepatosplenomegaly and only 20% patients with mild hepatosplenomegaly.
- 46% of thalassemic patients presented between -1SD & -2SD followed by 20% thalassemic patients presented between -3SD&-2SD followed by 20% thalassemic patients between less than <-3SD and only 14% thalassemic patients between -1SD & MEDIAN. Mean height was 105.77 cm (SD ± 13.79)

TABLE -1 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO HEMOGLOBIN LEVELS

AGE	MODREATE	SEVERE
<1YR	1(1.6%)	2(3.3%)
2-3YRS	6(10%)	7(11.6%)
4-6 YRS	2(3.3%)	12(20%)
7-9 YRS	4(6.6%)	3(5%)
>9 YRS	7(11.6%)	16(27%)
TOTAL	20(33%)	40(67%)

This table shows that 67% of thalassemic patient presented with severe anemia and 33% of thalassemic patients with moderate anemia. Mean hemoglobin level: 8.2±2gm/dl.




TABLE -2 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO BLOOD MCV LEVELS

AGE	MODERATE	SEVERE
<1YR	2(3.3%)	1(1.6%)
2-9YRS	9(15%)	27(45%)
>10YRS	10(16.6%)	11(18.5%)
TOTAL	21(35%)	39(65%)

This table shows that 65% of thalassemic patients had severe microcytic RBCs and only 35% of thalassemic patients had moderate microcytic RBCs. Mean MCV level: $73 \pm 2 \mu\text{m}^3$

TABLE -3 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO MCH LEVELS

AGE	MODERATE	SEVERE
<1YR	1(2%)	2(3.3%)
2-9YRS	10(16.6%)	26(43%)
>10YRS	9(15%)	12(20%)
TOTAL	20(24%)	40(66%)

This table shows that 66% of thalassemic patients had severe hypochromic RBCs. Mean MCH LEVEL: $23 \pm 2 \text{pg/cell}$

TABLE -4 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO BLOOD MCHC LEVELS

AGE	N=60
NORMAL(32-36)	3(5%)
ABNORMAL(HYPOCHROMIC)	57(95%)
TOTAL	60

This table shows that 95% of thalassemic patients had severe hypochromic blood. Mean MCHC level: $28 \pm 2 \text{g/dl}$

TABLE- -5 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO SERUM FERRITIN LEVELS

SERUM FERRITIN	N=60
<1000	8(12%)
1000-2500	29(49%)
>2500	23(39%)
TOTAL	60

This table shows that 49% of thalassemic patients had ferritin levels between 1000 -2500 followed by 39% of thalassemic patients more than 2500 levels and only 12% of thalassemic patients had lesser than 1000 levels. Mean ferritin level: $1281 \pm 219 \text{ng/dl}$

-55% of thalassemia patients had frequency of blood transfusion 5-10 times per year followed by 33% of patients between 10-15 times per year and only 12% patients more than 15 times per year. Mean frequency of blood transfusion 16 ± 2 times a year

TABLE-6 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO INTERVAL BETWEEN TRANSFUSION (DAYS)

INTERVAL BETWEEN TRANSFUSIONS(DAYS)	N=60
15-25 DAYS	44(73%)
25-35 DAYS	10(16%)
35-45 DAYS	6(11%)
TOTAL	60

In this study 73% of thalassemia patients were transfused blood between 15-25 days followed by 16% of patients were transfused between 25-35 days and only 11% of patients were transfused blood between 35-45 days. Mean interval between transfusion 22 ± 2 days .

TABLE -7 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO RANDOM BLOOD SUGAR LEVELS

BLOOD SUGAR	N=60
EUGLYCEMIA	52(86%)

HYPERGLYCEMIA	6(10%)
HYPOGLYCEMIA	2(4%)
TOTAL	60

This table shows that 10% Of thalassemic patients were presented with hyperglycemia ,4% of thalassemic patients with hypoglycemia and 86% of patients were euglycemia

TABLE -8 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO GLUCOSE TOLERANCE TEST

GLUCOSE TOLERANCE TEST(GTT)	N=60
IMPAIRED GLUCOSE TOLERANCE	4(6.5%)
DIABETIC	2(3.3%)
NORMAL	54(90%)
TOTAL	60

This stable shows that 90% of thalassemic patients had normal glucose tolerance test followed by 6.5% of thalassemic patients which had impaired glucose tolerance test .Only 3.3% of thalassemic patients had glucose tolerance test in diabetic range.

TABLE 9 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO SERUM CALCIUM LEVELS

SERUM CALCIUM	N=60
NORMAL	49(81%)
HYPOCALCEMIA	11(18%)
TOTAL	60

This table shows that 18% of thalassemic patients had hypocalcemia And mean serum calcium level:8.20±2mg/dl

TABLE 10 DISTRIBUTION OF THALASSEMIA PATIENTS ACCORDING TO SERUM THYROID LEVELS

LEVEL	N(60)
T4 & TSH (NORMAL)	56(93%)
T4 NORMAL & TSH INCREASE	1(1.6%)
T4 DECREASE & TSH INCREASE	3(4.4%)
TOTA;	60

This table shows that out of 60 thalassemic patients 93% of thalassemic patients had normal thyroid followed by 5.4% patients had subclinical hypothyroidism and only 1.6% thalassemic patient had overt hypothyroidism.

DISCUSSION

In present study out of 60 children 75% (n=45) were males and 25%(n=22) were females. It is comparable to the study done by Karimi et al. (2009)⁵⁵ in Iran done on 50 Thalassemia major children out of which 66% were (n=33) were males and 34% (n=17) were females. A similar study done by A. Azankeivan et al. (2008)⁴⁸ included 104 thalassemia major patients out of which 63% (n=75) were male and 37% (n=39) were female. In a study done by Ozyouk et al.(2015)⁶⁶ 61% were males and 39% were females out of the total 49 patient.

The prevalence of β- thalassemia trait varies between 3-17% because of consanguinity and caste and area endogamy. 4 Every year, ten thousand children with β-thalassemia major are born in India, which constitutes 10% of the total number in the world.5 HbE thalassemia is common in north-east parts of India.

In Present study group comprised of patients ranging between 6 month to 18 years with a mean age of 10.20 \pm 3 years. Similar observation M. Arora et al. study at Lady Harding Medical College, New Delhi, studied 30 children and 20 matched controls. The mean age of study group 11.83 \pm 1.91 years (range 9-17 years) and that of control group 12 \pm 2.17 years (range 9-17 years) was comparable. Arie Bacalo et al.(1992)¹⁹ in an Arab hospital studied 17 children aged 6-17years (average 11.2years). In F M Abu Ekteish et al.(2007)⁴⁷ studied study group had a range of 7.5-18 years with mean a mean age of 12.5(\pm 3.5 years).

Mean hemoglobin in present study was 6.244(\pm 1.66) gm/dl which is much lower than most of studies done. In study by Bijan Keikhaei et al. (2009)⁵³ at Ahwaz Jondishpour University of medical Sciences, Ahwaz, Iran mean hemoglobin was (8.5 \pm 0.9)gm/dl. In study by M.Arora et al. (2001)³⁰ found mean hemoglobin of patient was (9.87 \pm 1.45gm/dl). A similar study done by Kanj N. et al. (2000)²⁷ mean hemoglobin was (8.8 \pm 1.1)gm/dl. In a study done by Carnelli D'Angelo, Pecchiari et al.(2003)³³ mean hemoglobin was (9.3 \pm 1)gm/dl this reason behind such a low mean value of hemoglobin was because parents stay away from hospital, ignorance and literacy The pathogenesis of growth failure in thalassemia is multifactorial and is mainly due to transfusional iron overload and resulting endocrinopathies (GH deficiency , hypothyroidism, diabetes), nutritional deficiencies, and intensive use of chelating agents particularly deferoxamine. Other etiologies particularly in sub optimally treated children are increased metabolism, chronic anemia, and hypoxia . The anterior pituitary is particularly sensitive to iron associated free radical oxidative stress . Even a modest amount of iron deposition in anterior pituitary by MRI can interfere with its function. Dysregulation of the GH insulin like growth factors axis leads to growth hormone deficiency and growth deceleration. (Balgir RS et-al Hematology blood transfusion)²²

All children with Beta thalassemia major >10 yrs should undergo standing and sitting height every 6 months, bone age, growth hormone stimulation, insulin like growth factor(IGF)-1level,and IGF-BP3 level .

Frequent blood transfusion should be considered in patients with growth failure with reassessment for tapering or withdrawal when a sustained clinical benefit is achieved. (Weatherall DJ et-al ,Thalassemia a global health problem .)²³

In present study 50% of thalassemia patients presented with severe hepatosplenomegaly followed by 30% patients with moderate hepatosplenomegaly and only 20% patients with mild hepatosplenomegaly. Similar study done by Santosh Kumar et al (2016) at Department of Pediatrics, MGM Medical College and LSK Hospital, Kishanganj, Bihar found Out of 211 patients hepatomegaly was the most common clinical finding among the study population (57.8%), followed by splenomegaly (54.9%) .

In present study 67% of thalassemic patient presented with severe anemia and 33% of thalassemic patients with moderate anemia.

In present study 66% of thalassemic patients had severe hypochromic anemia and 34 % of thalassemic patients had moderate hypochromic anemia ,mean MCH value was 23 \pm 2 pg/cell

In present study 65% of thalassemic patients had severe microcytic RBCs and only 35% Of thalassemic patients had moderate microcytic RBCs, mean MCV value was 73 \pm 2 μ m³.

In present study 95% of thalassemic patients had severe hypochromic anemia and mean MCHC value was 28 \pm 2 gHb/dl.

Is characterized by reduced Hb level (<7 g/dl), mean corpuscular volume (MCV) > 50 <70 fl (femtoliters) and mean corpuscular Hb (MCH) >12<20 pg. In peripheral blood smear affected individuals show RBC morphologic changes (microcytosis, hypochromia cells) and nucleated RBC (i.e., erythroblasts). The number of erythroblasts is related to the degree of anemia and is markedly increased after splenectomy. Thalassemia intermedia is characterized by Hb level between 7 and 10 g/dl, MCV between 50 and 80 fl and MCH between 16 and 24 pg (Deyde VM et-al, Hematological disorder).²⁴

In present study 55% of thalassemia patients had frequency of blood transfusion 5-10 times per year followed by 33 % of patients between 10-15 times per year and only 12% patients more than 15 times per year. Similar study done by Neeraj shah et al(2010) at NHL Municipal Medical College, Ahmedabad found that 36% of thalassemic patients had blood transfusion more than 15times per year, 34% of thalassemic patients had blood transfusion between 5-10times per year and only 28% of thalassemic patients had transfused blood more than 10 times per year .

Severe anemia with hemoglobin <7 g% for more than 2 weeks is widely accepted as an indication to start blood transfusion.[13] The goal should be aimed to maintain a pre-transfusional Hb level of 9 to 10 g/dl and a post-transfusion Hb level of 13 to 14 g/dl. Such regime generally prevents growth impairment, organ damage and bone deformities. Care should be taken to avoid faster transfusion exceeding 5 ml/kg/h and amount of transfused RBC should not exceed 15 to 20 ml/kg/day. The frequency of transfusion is usually every 2 to 4 weeks (Ramesh agarawal et-al an overview of Thalassemia)²⁵.

In present study 49% of thalassemic patients had ferritin levels between 1000 -2500ng/ml followed by 39% of thalassemic patients more than 2500 level ng/ml and only 12% of thalassemic patients had ferritin level less than 1000 ng/ml. Similar study done by Anil.k. Saraya, M.D., et-al at Department of Pediatrics and Computer Center, ; AIIMS Delhi studied Serum ferritin levels in 64 patients with beta homozygous thalassemia (BHT), 120 patients with beta heterozygous thalassemia, and 46 normal subjects. Incidence of iron overload seen in 32 Beta heterozygous thalassemia cases was similar in untransfused and transfused cases. Among heterozygotes thalassemia patients, iron stores were depleted in 24 (20%) patients, mostly females 70.8% and 29.2% of males .

The iron burden on the body can be estimated by means of serum ferritin, iron, and TIBC levels. The estimation of serum ferritin levels is the most commonly employed test to evaluate iron overload in β -thalassemia major. The association between serum ferritin and levels of body iron are well-established and the test is easy to perform compared with other tests for iron overload. (Panchal and patel et-al Iron chelation status in young children with Thalassemia)²⁶

The conventional treatment at many centers for severe myocardial siderosis with heart failure is long-term, continuous, high-dose intravenous deferoxamine. Several small studies have confirmed that this approach is effective, and reversal of cardiomyopathy is possible

Combined chelation therapy in this situation might be effective, yet prospective trials examining the treatment of severe cardiac siderosis are lacking.

Since the body has no effective means of effectively removing iron, the only way to remove excess iron is to use iron chelators. The major step forward in improving survival and reducing complications was the introduction, of the chelating agent deferoxamine, used as a subcutaneous infusion. Two oral chelators, deferiprone and deferasirox, have recently become available, making therapy easier and more efficacious. Compliance, although improved by the switch to oral therapy, still presents a problem and is the major obstacle to effective prevention of iron overload. The orally

active chelators seem to be more effective in gaining access to the chelatable iron pool of cardiomyocytes, binding labile iron, and attenuating reactive oxygen species formation⁴⁶. Other study showed that the combination therapy is associated with lower risk of mortality

In present study, hyperglycemia was found in 10% of thalassemic patients and hypoglycemia in 4% of thalassemic patients and 86% of thalassemic patients were euglycemic. Using oral glucose tolerance test 90% of thalassemic patients had normal glucose tolerance test followed by 6.5% of thalassemic patients who had impaired glucose tolerance and only 3.3% of thalassemic patients had Glucose tolerance test in diabetic range. Similar study done by Swati Mohan Gadappa et al at Department of Paediatrics, Smt. Kashibai Navale Medical College and General Hospital, Pune.

In their study Found that out of 25 children 16% of thalassemic patients had impaired Glucose Tolerance and 84% of thalassemic patients had normal glucose tolerance. Similar study by Ashraf T. Soliman et al (2013) at Department of pediatrics Hamad Medical Centre Qatar found that out of 16 thalassemic children 25% patients had impaired fasting blood glucose concentration followed by 6.25% thalassemic patient had blood glucose level in diabetic range and 12.5% thalassemic patient had impaired glucose tolerance (IGT) (BG > 7.8 and <11.1 mmol/L).

Glucose intolerance in adolescence and diabetes mellitus later in life are also frequent complications mainly due to iron overload, chronic liver disease and genetic predisposition.¹⁵ The effect of iron overload on glucose metabolism is probably due to iron deposition in liver and pancreas causing increased insulin resistance or decreased insulin production leading to glucose intolerance and further diabetes mellitus.

In present study, out of 60 thalassemic patients 93% of thalassemic patients had normal thyroid function followed by 5% patients had subclinical hypothyroidism (normal T4 & high TSH) and only 1.6% thalassemic patients had overt hypothyroidism (high TSH & low T4). Similar study done by Rahul Panchal et-al at Government Medical College, Surat, Gujarat found that Thyroid function as indicated by the level of thyroid hormones was impaired in 10 out of 100 patients (10%); all 10 patients (10%) had subclinical hypothyroidism (normal T4 with high TSH); no case of secondary hypothyroidism (high TSH, T4) was found. Similar study done by Farideh Mogharabet-al (2017) Jahrom University of Medical Sciences, Jahrom, Iran found that Thyroid hormone level was normal in 106 of 112 total examined patients (94.6%) and only 6 patients (5.4%) suffered from hypothyroidism.

The most common form of thyroid dysfunction seen in thalassemia is primary hypothyroidism. Nonetheless, the frequency of hypothyroidism shows a discrepancy depending on the region, quality of management, and treatment protocols. Different studies suggest that thyroid dysfunctions appear with a frequency of 13%–60% in thalassemic patients after 10 years of age regardless of difference in the rate of prevalence, largely as in the form of subclinical hypothyroidism.²⁹

In present study 18% of thalassemic patients had hypocalcemia with mean serum calcium 8.3 ± 0.8 mg/dl. Similar study done by Salva Shah et al at Kabir Medical College, Peshawar - Pakistan found that frequency of hypocalcemia was 49%. Hyperphosphataemia was associated with 53% of hypocalcemic patients (Hypoparathyroidism was suspected in 26 patients). Mean serum calcium was 8.46 ± 0.94 mg/dl while mean phosphate level was 5.33 ± 0.77 mg/dl in the subjects.

Hypocalcemia occurring in β -thalassemia major is attributed to hypoparathyroidism mainly. Low parathyroid hormone levels lead to excessive calcium loss in urine, decrease bone remodeling and decreased intestinal absorption of calcium¹⁵. Hypocalcemia is a common biochemical abnormality which may be asymptomatic in mild cases or present as acute life-threatening crisis. Chronic hypocalcemia complications are mainly those of bone disease. In addition, severe hypocalcemia can result in cardiovascular collapse, hypotension which does not respond to fluids and dysrhythmias.

Neurological complications of low serum calcium are acute seizures, tetany, basal ganglia calcification, parkinsonism, hemiballismus and choreoathetosis.

CONCLUSION

Out of 60 patients of thalassemia were enrolled and their demographic hematological profile was taken. The mean age was 10.2 yrs, 75% were males and 81% were Hindus ,40% belonged to upper lower class .

Most common clinical features were (97%) icterus followed by (90%)pallor,80% of thalassemia patients had moderate to severe hepatomegaly. Mean height was 105.77 ± 14 cm ,Mean Hb was 8.5 ± 0.9 gm/dl, 67% had severe anemia , Mean MCV level was 73 ± 2 μ m³,Mean MCH and MCHC level were 20 ± 2 pg/cell, 28 ± 2 gHb/dl.

Mean ferritin level was 1281.8 ± 219.9 ng/dl , 45% of thalassemic patients had serum ferritin level between 1000 to 2000 ng/dl followed by 39% of patients had serum ferritin level more than 2500 ng/dl,Mean frequency of blood transfusion 16 ± 2 times a year , 55% of had blood transfusion 5-10 times a year followed by 33% had frequency of blood transfusion 10-15 times a year and only 12% had frequency more than 15 times a year , Mean interval between transfusion 22 ± 2 days ,73% had an interval of 15 to 25 days between two transfusion., 86% had euglycemia and 10% had hyperglycemia when blood sugar was tested randomly.

90% of thalassemic patients had normal glucose tolerance test followed by 6.5% of thalassemic patients had impaired glucose tolerance test,while 3.3 % patients came under diabetic range.

81% of thalassemic patients had normal serum calcium level while 18% had hypocalcemia,93% of thalassemic patients had normal serum T4,and TSH level ,1.6% of patients had subclinical hypothyroidism and only 4% of thalassemic patients had overt hypothyroidism.

Tablet deferasirox was taken as chelating agent in all thalassemia patients, 51% of which were taking it after 2years of age,35% were had started between 1-2 years of age and only 13% started below 1 year of age .

The common adverse reaction in thalassemic patients were diarrhea (26.1%), abdominal pain (23%),skin rash(20%),seizures(16%),blurring of vision (3%).

Hepatitis C virus infection was found in 76% and hepatitis B surface antigen (HBsAg) was positive in 48% of thalassemia patients.

Two patients (4%) out of 60 thalassemic patients had decreased left ventricular ejection fraction of 35 to 40%.

BIBLIOGRAPHY

1. Cooley, T. B. and Lec, P. (1925): A series of cases of splenomegaly in children with anemia and peculiar bone changes. *Trans. Am. Pediatr. Soc.* 37: 29.
2. Cooley, T. B. (1927): Von Jaksch's anemia. *Am. J. Dis. Child.* Chicago. 33: 786.
3. Bradford, W. L. and Dye, J. (1936): Observations on the morphology of the erythrocytes in disease thalassemia (Erythroblastic anemia of Cooley). *J. Pediatr.* 9: 3 12-13.
4. Cooley T B, M.D. 1871-1945. *Am J Dis Child* (1946); 71: 77 - 79.
5. Duke M, Abelmann WH. The hemodynamic response to chronic anemia. *Circulation* (1969);34:503–515.

6. Sukumaran, P.K. and Master (1973): The distribution of abnormal haemoglobins in the Indian population. In proceedings of the First Conference of the Indian Society of Human Genetics vol. 1: Human population genetics in India, Bombay. Orient Longman. 91-111.
7. Letsky EA, Miller F, Worwood M, et al. Serum ferritin in children with thalassaemia regularly transfused. *J Clin Pathol.* 1974;27:652–655.
8. Keens TG, O’Neal MH, Ortega JA, et al. Pulmonary function abnormalities in thalassemia patients on a hyper transfusion program. *Pediatrics.* (1980);65:1013– 1017.
9. Modell, B. and Petrou M. (1983): The problem of hemoglobinopathies in India. *Ind. J. Haematol.* 1-5.
10. JOHN C. WOODa et-al Physiology and Pathophysiology of Iron Cardiomyopathy in Thalassaemia
11. M. Baldini, I A. Marcon et-al Beta-Thalassaemia Intermedia: Evaluation of Endocrine and Bone Complications.
12. Addison, G. M., Beamish, M. R., Hales, C. N., Hodgkins, M., Jacobs, A., and Llewellyn, P. (1972). An immunoradiometric assay for ferritin in the serum of normal subjects and patients with iron deficiency and iron overload. *J. clin. Pathol.*, 25, 326-329.
13. Barry, M. (1973). Iron overload: clinical aspects evaluation and treatment. *Clin. Haematol.*, 2, no. 2.
14. Barry, M. (1974). Liver iron concentration, stainable iron, and total body storage iron. *Gut*, 15, 411-415.
15. Barry, M., Flynn, D. M., Letsky, E. A., and Risdon, R. A. (1974). Long-term chelation therapy in thalassaemia major: the effect on liver iron concentration, liver histology and clinical progress.
16. *Brit. med. J.*, 2, 16. Cappell, D. F., Hutchison, H. E., and Jowett, M. (1957). Transfusional complication in thalassemia children.
17. Weatherall DJ, Clegg JB. Thalassaemia Ñ a global public health problem. *Nat Med* 1996;2:847-9.
18. Higgs DR. *a*-Thalassaemia. *Baillieres Clin Haematol* 1993;6:117-50.
19. Chui DH, Waye JS. Hydrops fetalis caused by alpha-thalassemia: an emerging health care problem. *Blood* 1998;91:2213-22.
20. Weatherall DJ. The thalassemiias. In: Stamatoyannopoulos G, Nienhuis AW, Majerus PH, Varmus H, eds. *The molecular basis of blood diseases*. 2nd ed. Philadelphia: W.B. Saunders, 1994:157-205.
21. Flint J, Harding RM, Boyce AJ, Clegg JB. The population genetics of the haemoglobinopathies. *Baillieres Clin Haematol* 1993;6:215-62.
22. Weatherall DJ. Common genetic disorders of the red cell and the Òmalaria hypothesis. *Ó Ann Trop Med Parasitol* 1987;81:539-48.
23. Allen SJ, O’Donnell A, Alexander NDE, et al. *a*+Thalassaemia protects children against disease caused by other infections as well as malaria. *Proc Natl Acad Sci U S A* 1997;94:14736-41.
24. Anderson LJ, Westwood MA, Holden S, Davis B, Prescott E, Wonke B, Porter JB, Walker JM, Pennell DJ. 2004. Myocardial iron clearance during reversal of siderotic cardiomyopathy with intravenous desferrioxamine: A prospective study using T2_ cardiovascular magnetic resonance. *24Br J Haematol* 127: 348–355.
25. Bank A, Marks PA. 1966. Excess a chain synthesis relative to b chain synthesis in thalassaemia major and minor. *Nature* 212: 1198–1200.
26. Borgna-Pignatti C, Rugolotto S, De Stefano P, Zhao H, Cappellini MD, Del Vecchio GC, Romeo MA, Forni GL, Gamberini MR, Ghilardi R, et al. 2004. Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica* 89: 1187–1193.

27. Borgna-Pignatti C, Cappellini MD, De Stefano P, Del Vecchio GC, Fomi GL, Gamberini MR, Ghilard R, Piga A, Romeo MA, Zhao H, et al. 2006. Cardiac morbidity and mortality in deferoxamine- or deferipronetreated patients with thalassemia major. *Blood* 107: 3733–3737.
28. Borgna-Pignatti C, Marsella M, Zanforrin N. 2010. The natural history of thalassemia intermedia. *Ann NY Acad Sci* 1202: 214–220.
29. Boyer Sh, Belding TK, Margolet L, Noyes AN. 1975. Fetal hemoglobin restriction to a few erythrocytes (F cells) in normal human adults. *Science* 188: 361–363.
30. Breuer W, Ermers MJ, Pootrakul P, Abramov A, Hershko C, Cabantichik ZI. 2001. Desferrioxamine-chelatable iron, a component of serum non-transferrin-bound iron, used for assessing chelation therapy. *Blood* 197: 792–798.
31. Brittenham GM, Farrell DE, Harris JW, Feldman ES, Danish EH, Muir WA, Tripp JH, Bellon EM. 1982. Magnetic- susceptibility measurement of human iron stores. *N Engl J Med* 307: 1671–1675.
32. Brittenham GM, Cohen AR, McLaren Ce, Martin MB, Griffith PM, Nienhuis AW, Young NS, Allen CJ, 31.Farrell DE, Harris JW. 1993. Hepatic iron stores and plasma ferritin concentration in patients with sickle cell anemia and thalassemia major. *Am J Hematol* 42: 81–85.
33. Brittenham GM, Griffith PM, Nienhuis AW, McLaren CE, Young NS, Tucker EE, Allen CJ, Farrell DE, Harris JW. 1994. Efficacy of deferoxamine in preventing complications of iron overload in patients with thalassemia major. *N Engl J Med* 1331: 567–573.
34. Brittenham GM, Nathan DG, Olivieri NF, Pippard MJ, Weatherall DJ. 2003. Deferiprone versus desferrioxamine in thalassaemia and T2_ validation and utility. *Lancet* 361: 183.
35. Cao A, Galanello R. 2010. b-Thalassemia. *Genet Med* 12: 61–76.
36. Cao A, Galanello R, Rosatelli MC. 1994. Genotype–phenotype correlations in b-thalassemias. *Blood Rev* 8: 1–12.
37. Cappellini MD, Motta I, Musallam KM, Taher AT. 2010. Redefining thalassemia as a hypercoagulable state. *Ann NY Acad Sci* 1202: 231–236.
38. Weatherall D J and Clegg J B. *The thalassaemia syndromes*. 4th ed. Oxford, England: Blackwell Scientific (in press). Cited by Olivieri N F. *THE β -THALASSEMIAS*. *The New England Journal of Medicine*, (1999); 341(2): 99-109.
39. Camaschella C, Cappellini MD. *Thalassemia intermedia*. *Haematologica*, (1995); 80: 58-68. PMID:7758995.
40. Cao A. *Diagnosis of b-thalassemia intermediate presentation*. In: Fucharoen S, Rowley PT, Paul NW, eds. *Thalassemia: pathophysiology and management*. Part B. Vol. 23 of *Birth defects: original article series*. New York: Alan R. Liss, (1988): 219-26.
41. Ho PJ, Hall GW, Luo LY, Weatherall DJ, Thein SL. *Beta thalassaemia intermedia: is it possible consistently to predict phenotype from genotype?* *Br. J. Haematol.* (1998); 100: 70-8. PMID:9450794.
42. Giardine B, van Baal S, Kaimakis P, Riemer C, Miller W, Samara M, Kollia P, Anagnou NP, Chui DH, Wajcman H, Hardison RC, Patrinos GP: *HbVar database of human hemoglobin variants and thalassemia mutations: 2007 update*. *Hum. Mutat.* (2007), 28: 206. PMID:17221864.
43. Huisman THJ, Carver MFH, Baysal E: *A Syllabus of Thalassemia Mutations*. The Sickle Cell Anemia Foundation, Augusta, GA (1997).
44. Galanello R, Cao A: *Relationship between genotype and phenotype*. *Thalassemia intermedia*. *Ann. N Y Acad. Sci.* (1998), 850: 325-333. PMID: 9668554.
45. Uda M, Galanello R, Sanna S, Lettre G, Sankaran VG, Chen W, Usala G, Busonero F, Maschio A, Albai G, et al. *Genome-wide association study shows BCL11A associated with persistent fetal hemoglobin and amelioration of the phenotype of beta-thalassemia*. *Proc. Natl. Acad. Sci. USA.* (2008); 105: 1620-1625.

46. Origa R, Satta S, Matta G, Galanello R: Glutathione S-transferase gene polymorphism and cardiac iron overload in thalassemia major. *Br. J. Haematol.* (2008); 142: 143-145. doi: 10.1111/j.1365-2141.2008.07175.
47. Sollaino MC, Paglietti ME, Perseu L, Giagu N, Loi D, Galanello R: Association of alpha globin gene quadruplication and heterozygous beta thalassemia in patients with thalassemia intermedia. *Haematologica*, (2009); 94: 1445-1448. doi: 10.3324/haematol.2009.005728.
48. Taher A T, Musallam K M and Karimi M. et al., “Overview on practices in thalassemia intermedia management aiming for lowering complication rates across a region of endemicity: the optimal care study,” *Blood*, (2010); 115 (10): 1886–1892. doi: 10.1182/blood-2009-09-243154. Epub 2009 Dec 23.
49. Galanello R and Origa R. Beta thalassaemia. *Orphanet J. Rare Diseases*, (2010); 5 (11): 21. doi:10.1186/1750-1172-5-11.
50. A. Haddad A. , Tyan P., Radwan A., Mallat N., and Taher A. “ β - thalassemia intermedia: a bird’s-eye view,” *Turkish Journal of Hematology*, (2014); 31 (1): 5. doi: 10.4274/Tjh.2014.0032.