

A CLINICAL STUDY ON TESTICULAR TORSION IN SVMC TIRUPATHI

- 1. Dr. G. Kavitha, Assistant Professor of Surgery, SVMC, Tirupati**
- 2. Dr. P. Gowthami, Assistant Professor of Surgery, SVMC, Tirupati**
- 3. Dr. P. Sabitha, Associate Professor of Surgery, SVMC, Tirupati**

Corresponding Author: Dr. P. Sabitha, Associate Professor of Surgery, SVMC, Tirupati

ABSTRACT:

Introduction: Testicular torsion refers to the torsion of the spermatic cord structures and subsequent loss of the blood supply to the ipsilateral testicle. This is a surgical emergency; early diagnosis and treatment are vital to save the testicle and preserving future fertility.¹ The rate of testicular viability decreases significantly after 6 hours from onset of symptoms.² Testicular torsion is primarily a disease of adolescents and neonates. It is the most common cause of testicular loss in these age groups. However, torsion may occasionally occur in men 40-50 years old.³ **Aim** of the study is to know the clinical presentation, management of diagnosed cases of testicular torsion. **Methodology:** An prospective observational study done in 13 cases diagnosed to have Testicular Torsion from January 2023 to August 2023 in the Department of General Surgery, Sri Venkateswara Medical College, Tirupati. History, clinical examination findings, investigations, management and complications were noted in the proforma and the results were analyzed for the study purpose. **Results:** In the present study majority of the cases were in the age group of 31-40 years. All cases had scrotal pain and cases with delayed presentation had fever and swelling. Only 2 cases presented before a day and 54% cases presented between 1- 3 days and 31% of cases presented more than 3 days. **Conclusion:** Testicular torsion is a acute surgical emergency

and needs immediate intervention to prevent loss of testis and fertility. So all acute scrotal conditions should be considered as torsion and should be evaluated by USG Doppler to rule out torsion. Orchidopexy of the contra lateral testis is needed to prevent torsion in the contra lateral side.

KEYWORDS: Testicular Torsion, Orchicedctomy, Orchidopexy

Introduction:

Testicular torsion refers to the torsion of the spermatic cord structures and subsequent loss of the blood supply to the ipsilateral testicle. This is a surgical emergency; early diagnosis and treatment are vital to save the testicle and preserving future fertility.¹

The rate of testicular viability decreases significantly after 6 hours from onset of symptoms.² Testicular torsion is primarily a disease of adolescents and neonates. It is the most common cause of testicular loss in these age groups.

However, torsion may occasionally occur in men 40-50 year's old.³ Surgical treatment may prevent further ischemic damage to the testis. Diagnosis of testicular torsion is clinical, and diagnostic testing should not delay treatment.

Torsion can be intravaginal torsion and extravaginal torsion. Causes of Testicular torsion can be due to anomalies like bell clapper deformity, trauma, cryptorchidism, sudden contraction of the cremasteric muscles, testicular malignancy⁵. Occurs in about 17% of males⁴ and is bilateral in 40%.

Torsion should be differentiated from epididymo orchitis, testicular hematoma post trauma, complications of hydrocele etc. Ultrasound with Doppler will help in the differentiation.

The TWIST (Testicular Workup for Ischemia and Suspected Torsion) scoring system was developed for the purpose of determining the risk of testicular torsion on clinical grounds, thus decreasing the indication for ultrasound.⁶

Ultrasonographic findings suggestive of acute testicular torsion include the following⁷:

Absent or decreased blood flow in the affected testicle, Decreased flow velocity in the intratesticular arteries, Increased resistive indices in the intratesticular arteries, Hypervascularity with a low resistance flow pattern (after partial torsion-detorsion). Effective diagnosis and intervention at the early stage will prevent or help in avoiding orchidectomy.

Aim of the study: To study the clinical presentation, management of diagnosed cases of testicular torsion.

Methodology: A prospective observational study done in 13 cases diagnosed to have Testicular Torsion from January 2023 to August 2023 in the Department of General Surgery, Sri Venkateswara Medical College, Tirupati. History, clinical examination findings, investigations, management and complications were noted in the proforma and the results were analyzed for the study purpose.

Observations and results: During the study period about 13 cases were studied.

Table no 1: Age distribution of the cases

Age in years	No of cases	Percentage
11 - 20	4	31%
21-30	2	15%
31-40	2	15%
41-50	5	38%
Total	13	100

Graph no 1: Age distribution of the cases




Table no 2: Clinical Presentation

Presentation	No of cases	Percentage
Scrotal pain	13	100
Swelling	7	47
Fever	4	27

Graph no 2: Clinical Presentation




Table no 3: Duration of symptoms

Duration	No of cases	Percentage
<1day	2	15%
1-3 days	7	54%
>3days	4	31%

Graph no 3: Duration of symptoms





Table no 4: Management of cases

Management	No of cases	Percentage
Orchidectomy with orchidopexy	11	85%
Orchidopexy alone	2	15%

Graph no 4: Management of cases



Discussion:

In the present study majority of the cases were in the age group of 41- 50 years. All cases had scrotal pain and cases with delayed presentation had fever and swelling. Only 2 cases presented before a day and 54% cases presented between 1- 3 days and 31% of cases presented more than 3 days.

Only 2 cases had early diagnosis and had bilateral Orchidopexy where as the remaining cases who presented late were treated outside by non medical persons conservatively and presented with complications like testicular gangrene and abscess and had orchidectomy on the affected side and Orchidopexy on the contra lateral side.

Chu and colleagues noted that postoperative viability occurred in 95% (19/20) versus 67% (8/12) of patients with ischemia times of 24 hours or less and more than 24 hours,

respectively. Atrophy occurred in 67% (12/18) versus 83% (10/12) of these groups, respectively.⁸

Grimsby and colleagues prospectively studied 56 children with torsion to determine predictors of atrophy. All underwent an Orchidopexy irrespective of intraoperative findings. Grimsby and colleagues found that a black or hemorrhagic testis 5 minutes after detorsion, pain duration of longer than 12 hours, and heterogeneous parenchyma on preoperative US were associated with significant testis volume loss on follow-up US compared with a normal testis. All patients with a black or hemorrhagic testis had more than 80% volume loss.⁹

Conclusion: Testicular Torsion is an acute surgical emergency and needs immediate intervention to prevent loss of testis and fertility. So all acute scrotal conditions should be considered as torsion and should be evaluated by USG Doppler to rule out torsion. Orchidopexy of the contra lateral testis is needed to prevent torsion in the contra lateral side.

References:

1. Ta A, D'Arcy FT, Hoag N, D'Arcy JP, Lawrentschuk N. Testicular torsion and the acute scrotum: current emergency management. *Eur J Emerg Med.* 2015 Aug 11. 37-41.
2. Barbosa JA, Denes FT, Nguyen HT. Testicular Torsion-Can We Improve the Management of Acute Scrotum?. *J Urol.* 2016 Jun. 195 (6):1650-1.
3. Acute Scrotum. American Urological Association. July 2016; Accessed: November 22, 2016.
4. Barthold JS. Abnormalities of the Testis and Scrotum and their Surgical Management. McDougal WS, Kavoussi LR, Novick AC, Partin AW, Peters CA, eds. *Campbell- Walsh Urology.* 10th ed. Philadelphia, PA: WB Saunders; 2012. 642-5.
5. Uguz S, Yilmaz S, Guragac A, Topuz B, Aydur E. Association of Torsion With Testicular Cancer: A Retrospective Study. *Clin Genitourin Cancer.* 2016 Feb. 14 (1):e55-7.

6. Barbosa JA, Tiseo BC, Barayan GA, Rosman BM, Torricelli FC, Passerotti CC, et al. Development and initial validation of a scoring system to diagnose testicular torsion in children. *J Urol*. 2013 May. 189 (5):1859-64.
7. Prando D. Torsion of the spermatic cord: the main gray-scale and doppler sonographic signs. *Abdom Imaging*. 2009 Sep-Oct. 34(5):648-61.
8. Chu DI, Gupta K, Kawal T, et al. Tunica vaginalis flap for salvaging testicular torsion: a matched cohort analysis. *J Pediatr Urol*. 2018 Feb 6. pii: S14775131(18)30037-8.
9. Grimsby GM, Schlomer BJ, Menon VS, et al. Prospective evaluation of predictors of testis atrophy after surgery for testis torsion in children. *Urology*. 2018;116:150-155.