

INTRAOPERATIVE COMPARISON BETWEEN ISOFLURANE AND NITROGLYCERIN FOR HYPOTENSIVE ANAESTHESIA

Dr Nirod kumar Sahoo¹, Dr Nagendra Kumar Rajsamant², Dr Sujit Kumar Mohanty³, Dr Mrutyunjaya Mallick⁴

¹Asst. Professor, Dept of General Surgery , MKCG Medical College , Brahmapur, Ganjam, Odisha, India.

Mail: dr.nirodsahoo@gmail.com

²Assistant Professor Dept of Surgery, SCB MCH, Cuttack, Odisha, India

³Asst Professor, Dept of General Surgery , SCBMCH, Cuttack, Odisha, India

⁴Asst. professor, Dept of Anesthesiology & Critical care, MKCG Medical College , Brahmapur, Ganjam, Odisha, India

Corresponding author: Dr Mrutyunjaya Mallick, Asst. Professor, Dept of Anesthesiology & Critical care, MKCG Medical College, Brahmapur, Ganjam, Odisha, India. Mail:

mallick.drmrutyunjaya@gmail.com

Abstract

Introduction

All surgeries involve cutting of the blood vessels which will obviously result in bleeding.

Deliberate hypotension is a method to produce a controlled and safe reduction in the intravascular pressure, while preserving organ perfusion.

By enhancing the visualization of the surgical field, hypotension allows accurate delineation of lesions thereby causing fewer traumas to the delicate structures. Therefore intricate operations may be performed easily, exactly and therefore more successfully. Extensive surgical procedures are associated with considerable haemorrhage during and after surgery.

The concept of intentionally decreasing arterial BP to hypotensive levels during surgery was first proposed by **Cushing** in **1917** for intracranial surgery and was introduced into clinical practice by **Gardner** in **1946**.

Since then, this technique has been variously named as **controlled hypotension, Induced hypotension, deliberate hypotension and hypotensive anaesthesia**.

Deliberate hypotension is defined as the intentional reduction of the systemic perfusion pressure. Most studies define deliberate hypotension as reduction in systolic blood pressure (SBP) to 80-90mm Hg (30% decrease in the SBP from the baseline pressure). According to another definition, deliberate hypotension is a decrease in the mean arterial pressure (MAP) to 50-65 mm Hg in normotensive patients.

Main purpose of hypotensive anaesthesia is to decrease intraoperative blood loss, thereby improving operating condition or decreasing the need for blood transfusions.

Different techniques have been used to achieve controlled hypotension.

1. PHYSIOLOGIC TECHNIQUES:-

Body positioning, the haemodynamic effects of mechanical ventilation, and changes in heart rate and circulatory volume can be used with drugs to lower blood pressure to the desired level. The

appropriate use of physiologic maneuvers helps decrease the dose of potentially toxic drugs needed to produce hypotension.

2. PHARMACOLOGIC TECHNIQUES:- These includes:-

- Volatile Anaesthetics -- Halothane, Isoflurane, Sevoflurane, Enflurane, Desflurane.
- Directly acting vasodilators -- Sodium Nitroprusside, Nitroglycerine, Hydralazine, Adenosine.
- Ganglion Blocking Agents -- Trimethapan
- Beta Blockers -- Propranolol, Esmolol

13

- Combined α & β blocker -- Labetalol
- Calcium channel blocker -- Nicardipine
- Prostaglandins -- PGE1
- Spinal and Epidural Anaesthesia

The **ideal hypotensive agent** should have a predictable, dose dependent effect, rapid onset and recovery, quick elimination without producing toxic metabolites and should not disrupt auto regulation of vital organs.

Nitroglycerine is an organic nitrate which causes direct nonspecific, vascular smooth muscle relaxation, consequently decreasing both preload and after load on heart leading to fall in both systolic and diastolic blood pressure.

Isoflurane is a halogenated methyl ether and a potent inhalational anaesthetic. Isoflurane produces rapid hypotension by decreasing peripheral vascular resistance without myocardial depression . It also provides better myocardial perfusion than halothane or enflurane . With isoflurane baroreflex activity is better maintained in hypotensive anaesthesia .It 14 has rapid onset and recovery , more potent peripheral vasodilating effect .It also has the ability to protect the brain from hypoxia due to decrease in cerebral metabolic rate . (**Artru 1984 : Michenfelder et al in 1987**). With these advantages , Isoflurane has been gaining increasing interest recently as a hypotensive agent in induced hypotension . Isoflurane has been used as the sole agent to induce hypotension in orthoganthic surgery and has significantly reduced blood loss and improved the quality of surgical field .

Since both these drugs are safe and easily available , an attempt has been made to study comparing Nitroglycerine with Isoflurane as hypotensive agents during spinal surgeries in the Operation theatres of SCB Medical College and Hospital ,Cuttack.

Materials & Methods

The study titled “ **Comparison between Nitroglycerine and Isoflurane for Hypotensive anaesthesia** ” was carried out at SCB Medical College and Hospital ,Cuttack, in patients of both the sexes in the age group of 18 – 60 years belonging to ASA (American Society of Anaesthesiologists) grade I and II posted for elective spine surgery (lumber laminectomy and decompression) in operation theatres under general anaesthesia . The study was undertaken during the period August 2014 to July 2015.

After getting due permission from the Hospital ethical committee , informed consent was obtained from sixty patients selected

at random . They were thoroughly assessed preoperatively to exclude the presence of any significant systemic illness other than the disease for which they have taken admission .

The cases suitable for the study were prepared for surgery .

PREOPERATIVE ASSESSMENT : ---

Preoperative assessment was done by --

History .

74

Clinical examination .

Laboratory investigations .

1. HISTORY : --

Patients having past history of

- Central Nervous system and cardio vascular system diseases like Cerebro vascular accidents , Transient Ischemic Attacks , Myocardial Infarction , Angina or Intermittent Claudications .
- Brain pathologies like h/o migraine , tumor or epilepsy .
- Asthma , breathlessness on exertion or coagulation disorders .
- Drug therapy such as β –blockers , oral contraceptive pills , antihypertensives or anti –coagulant therapy .
- Liver and kidney dysfunctions .
- Relevant endocrinological disorders .
- Addiction to alcohol , smoking and narcotics .
- Grossly anaemic and hypovolemic patients .
- Chronically diseased and debilitated patients.

Were excluded from the study .

75

2. CLINICAL EXAMINATION : --

This was done by :-

- i. General Examination
- ii. Systemic Examination
- iii. Specific Examination

i) General Examination:

A detailed physical examination was done and patients having gross pallor , icterus , cyanosis , clubbing , oedema , lymphadenopathy , raised JVP and ascites were excluded from the study .

ii) Systemic Examination:

All the systems were examined thoroughly in every patient and if any abnormality detected then excluded from the study .

iii) Specific Examination:

To assess :

- The upper airway patency by inspecting nose , buccal cavity and pharynx .

76

- Difficult intubations by Mallampati Classification , patients

having more than grade II were excluded .

- Cardio respiratory reserve by sabrasez test , breath holding for more than 30 seconds .

- Overall fitness for general anaesthesia .

3) LABORATORY INVESTIGATIONS : -

i) Routine Tests

- Haemogram—Hb%

- DC ,TLC ,TRBC

- BT , CT , Blood Sugar (fasting and 2 hrs post prandial)

- Serum Urea , Creatinine for renal function assessment .

- Serum Electrolytes (sodium and potassium).

- Platelet count .

Urine :

- Routine

- Microscopic

Stool :

77

- Routine

- Microscopic

- ECG and CXR – Routinely in patients over 40 yrs of age and in cardio respiratory abnormality .

If any abnormality detected the cases were excluded from the study .

ii) Special :

The following special investigations were done when felt necessary and if any abnormality detected than the patients were excluded from the study .

- Liver function tests

- Lung function test

- 2D – ECHO

- X –Ray skull , CT Scan , EEG

- Ultrasono Gram of abdomen and renal system .

- Bllod urea , serum creatinine

- Serum hormone assays (e.g. thyroid)

78

PREOPERATIVE PREPARATION :

After thorough assessment and confirmed diagnosis , sixty patients were thus selected and all the cases were kept over night fasting or 6hr before surgery and tablet ranitidine 150mg and tablet Alprazolam 0.5mg was given orally on the night before operation.

Patients were then divided into two demographically identical groups of 30 each at random and informed consent for the study was taken .

They were named into two groups depending upon the type of hypotensive agent used .

□ Group N: Patients received IV Nitroglycerine infusion.

□ Group I: Patients received inhalation Isoflurane by fluotec vaporizer.

On the of operation patients were brought to the preanaesthetic room

atleast 60 minutes before surgery in the early hours of morning (around 9 .00 AM) and one intravenous infusion was started with 18 G venous cannula . Intravenous infusion of Ringers lactate solution was administered at the rate of 8 – 10 ml/kg/hr to maintain hourly urine output of atleast 1ml/kg/hr (50 ml or more).

In the operation room a preoperative check of blood pressure , pulse rate , SPO2 was done and ECG cables were attached for monitoring .

79

Premedication :

They were premedicated with inj. Glycopyrrolate 0.2mg i.v., inj. Butrophanol 0.2mg/kg i.v. and injection Midazolam 0.05mg/kg i.v. 20-30 mins prior to induction. Patients will be preoxygenated for 3 minutes with 100% oxygen at a flow rate of 5lit/min.

Induction :

All the patients were induced with an induction dose of Propofol 2.5mg/kg intravenously .

Intubation :

All the patients were intubated with appropriate size cuffed endotracheal tube after adequate relaxation by vecuronium bromide 0.1mg/kg. and mask ventilation with 100% oxygen . The cuff was inflated just to obliterate audible leakage , and fixed with adhesive tape .

Maintainance :

Anesthesia will be maintained with 60% N2O in oxygen; adequate muscle relaxation will be achieved with intermittent vecuronium bromide. Ventilation (tidal volume 8-10 ml/kg) will be adjusted to maintain end tidal carbon dioxide <35 mmHg.

80

Under all aseptic precautions a 15 cm central venous catheter will be inserted via the right internaljugular vein. Radial artery was cannulated using 20 G cannula , connected to a transducer for beat by beat blood pressure monitoring . All the patients will be catheterized with Foley's catheter for measuring urine output.

As surgery will be performed in prone position firm supports under chest and pelvis will be kept so that the abdominal movements and the venous return will not hampered. Compression of the abdomen by faulty positioning would result in the increase in central venous pressure (CVP) and engorged epidural veins. The eyes will be closed and covered and arms will be padded.

The selected hypotensive agent will be started after changing the patient to prone position.

The group N received Nitroglycerine infusion by syringe pump which was prepared by addition of 25 mg of nitroglycerine to 50 ml of 0.9% NS . The infusion was started with 1µg/kg/min and titrated to achieve the systolic blood pressure to less than 100 mm Hg before the skin incision and maintained in the range of 80 – 100 mm Hg or maintained the mean arterial pressure between 60 – 70 mm Hg . However it was decided not to exceed

the maximal rate of Nitroglycerine infusion above 10 µg/kg/min .

81

In group I isoflurane will be started at 0.5 vol % and adjusted to achieve the systolic blood pressure to less than 100 mm Hg before the skin incision and maintained in the range of 80 – 100 mm Hg or maintained the mean arterial pressure between 60 – 70 mm Hg . However it was decided not to exceed the inspired concentration of Isoflurane above 4% .

Monitoring of following parameters will be done: BP, HR, SpO₂, ETCO₂, ECG, CVP, Temperature, urine output.

The systolic blood pressure was lowered to less than 100 mm Hg before the skin incision and maintained in the range of 80 – 100 mm Hg throughout the surgery in both the nitroglycerine (Group – N) and Isoflurane (Group – I) groups , by adjusting the dose of nitroglycerine and concentration of Isoflurane . The blood loss was replaced by whole blood when the loss exceeds 10% Of estimated blood volume .Surgens were advised to not to use any adrenaline infiltration before the skin incision .All datas was recorded on a specially prepared anaesthesia record sheet .

The loss of blood was assessed by weighing of tetras , blood in the suction bottle and in the kidney tray deducing any amount of irrigating fluid used during the surgery .The swabs were weighted before they were used and weighted again after they were soaked with blood and thrown

82

individually into a collecting basket . The difference in the weight was taken as the amount of blood loss . 1gm gain in weight = 1 ml of blood loss .

For evaluation of the visibility of the operative field during surgery , the quality scale proposed by **Fromm and Boezaart** was used .

Grade – 1 – Slight bleeding – No suctioning of blood required .

Grade – 2 -- Slight bleeding – Occasional suctioning required . Surgical field not threatened .

Grade – 3 – Slight bleeding – Frequent suctioning required . Bleeding threatens surgical field a few seconds after suction is removed .

Grade – 4 – Moderate bleeding – Frequent suctioning required . Bleeding threatens surgical field directly after suction is removed .

Grade -5 -- Sever bleeding – Constant suctioning required . Bleeding appears faster than can be removed by suction . Surgical field threatens severely and surgery impossible .

The hypotensive agents were discontinued before suturing of the wounds.

83

The systolic blood pressure was allowed to return to pre hypotensive levels or near to it to check for the hemostasis .

The operative time was measured from the start of skin incision to the end of skin closure .

Fluid input during surgery period will be determined based on preoperative fasting, blood loss and clinical criteria (arterial pressure, heart rate and observation of the patient).

Ondansetron 8 mg i.v. was given before end of the surgery .

Reversal :

The patients were reversed at the end of surgery with appropriate doses of inj. Neostigmine (0.05 mg /kg) and inj. Glycopyrolate (0.01 mg/kg) Then the patients were kept in recovery room till they fully recovered . Then the patients were shifted to the ward .

Monitoring :

The patients were constantly monitored for vital functions like : NIBP, HR, SpO₂, ETCO₂, ECG, CVP, Temperature, urine output from premedication till the recovery and recorded as follows : -

(a) Before premedication

84

(b) Before induction,

(c) Before intubation ,

(d) After intubation ,

(e) Every 5 minutes till 30 minutes and then 10 minutes till the end of surgery .

Post operatively the patients were monitored in ICU and the above parameters were noted every two hourly .

If wide and frequent variations in the vital functions were observed they were recorded and steps were taken to correct them . These patients were excluded from study .

SPECIFIC OBSERVATIONS :

Pulse , blood pressure , ECG recording during and before induction of hypotension was observed .

(i) Pre operatively .

(ii) Before premedication .

(iii) Before intubation .

(iv) At the start of hypotensive agents .

85

(v) 5 minutes after starting the hypotensive agent and then every 5 minutes interval for 30 mins and then at 10 mins interval till the end of surgery .

(vi) at hypotensive agent discontinuation

(vii) after extubation

(viii) Blood loss was assessed from the suction bottle , weight difference of dry and wet tetra and from the kidney tray by a 20 cc syringe .

At the end of surgery dizziness , nausea , drowsiness , or any other postoperative complications were assessed .

The following points were specifically noted during induced hypotension in both the groups .

(i) Speed of onset of hypotension :

The time taken to lower systolic blood pressure to less than 100 mmHg or MAP to less than 70 mm Hg was noted in both the groups .This was taken as induction time .

(ii) Heart rate variations during induced hypotension.

(iii) Speed of recovery from hypotension .

86

The time taken for spontaneous recovery of systolic blood pressure to pre – induction value after discontinuation of the hypotensive agent was observed in both the groups was taken as the recovery from hypotension .

(iv) Effectivity of the hypotensive technique :

This was judged , taking into account the following parameters :

- (a) Mean intra –operative blood loss
- (b) Duration of surgery
- (c) No. of patients requiring intra –operative blood transfusion .
- (d) Surgeons score (about the dryness of surgical field)
- (e) Incidence of post –operative complication.

The data were entered into a Microsoft Excel Computer programme . Data were analysed using Medcal software . Descriptive statistics in the form of frequencies , Mean ,Median, Standard Deviations and was compared by using unpaired T –Test and data in proportion was compared by using Chi –square Test and Fisher’s exact test ($p < 0.05 =$ significant .).

Observations

A clinical comparative study of the use of Glyceryl Trinitrate (GTN) and Isoflurane in controlled intraoperative hypotension during spine surgery was undertaken in sixty patients of ASA Gr. I and II Of either sex , varying from 18 – 60 yrs of age , at S.C.B. Medical college and Hospital , Cuttack .

The observations were done as per the following parameters .

- 1. Age.
- 2. Sex .
- 3. Body weight.
- 4. Cardiovascular changes with respect to ;
 - Heart rate
 - Systolic Blood Pressure
 - Diastolic Blood Pressure
 - Mean Arterial Pressure.
- 5. Speed of onset of Hypotension in both the groups
- 6. Speed of recovery of hypotension in both the groups
- 7. Blood loss
- 8. Surgeon’s score
- 9. Percent fluctuation of various parameters.

HR and BP were recorded before premedications , before intubation , at the time of start of hypotensive agent and 1 , 5 , minutes after administration of hypotensive agent and then every 5 minutes till 30 minutes and then every 10 minutes till the end of surgery . Post reversal half hourly during recovery then two hourly in the recovery room . Again on the next day morning.

Discussion

The study “ **The comparison of Nitroglycerine and Isoflurane in Hypotensive anaesthesia** ” is carried out in spinal surgery procedures in S.C.B Medical college and Hospital , Cuttack taking sixty healthy ASA grade I and II patients posted for elective surgeries on spine (Lumber Laminectomy and Decompression) belonging to both sexes , and in the group of 18 – 60 years . The study was carried out during the period of august 2014 to July 2015 .

The result of the study are discussed with observations of other workers in the field of work , taking steps to account for difference or similarity as far as possible . The various aspects of discussed this study are as follows : --

AGE OF THE PATIENTS : --

In the Nitroglycerine group (Group – N) and Isoflurane group (Group - I) , the mean age of the patients are 39.300 ± 6.482 and 39.00 ± 7.86 years respectively . Almost equal age wise distributions are observed in both the groups . The age group 18 – 60 yrs was chosen to circumvent the variables at the extremes of age . We have deliberately excluded the geriatric age group because of the associated changes in cardiovascular physiology which would have interfered with our study . There is decreased

105
arterial compliance , decreased beta adrenergic responsiveness , decreased baroreceptor sensitivity , decreased SA node automaticity in geriatric patients . (**Harrisons Internal Medicine 2011, 19 th Edn.**) . Similarly younger patients are more resistant to the action of hypotensive drugs because of an exaggerated baro receptor response . They too were excluded from the study for this reason .

SEX OF THE PATIENTS : --

In Group – N and Group – I the male female ratio was respectively . Almost equal sex wise distribution are observed in both the groups . In our study too there was a male preponderance 50(83.33%) and female 10 (16.66%) . This male preponderance can be attributed to the fact that Men are devoted to the farming activities and are laborers lifting heavy weights, hence many workmen engage in activity that involves strain or trauma of lumbar spine.

WEIGHT OF THE PATIENTS : --

The weight of the patients had been considered for the purpose of dosages calculations of different types drugs used . Patients in both the

106
Groups didn't show any significant variations and were in the range of 35 – 70 kg.

Thus all the patients were demographically identical in relation to age , sex , and weight in both the groups .

Table No. – IV shows the preoperative vitals in the two study groups .
Heart rate , SBP , DBP , MAP were in the identical range in the two groups .

ONSET OF HYPOTENSION : --

In the present series , time of onset of hypotension was measured as the time to lower SBP to 100 mm Hg or below .Our study differs from the work of **Pal and Parmink (1976)** who judged the onset of action as the time to lower the SBP below 80 mm Hg .

We have maintained SBP in the range of 80 – 100 mm Hg and the MAP pressure around 70 mm Hg in order to prevent the disruption of autoregulatory blood flow to vital organs .

Nitroglycerine could reduce SBP faster than Isoflurane . In our study the mean onset of hypotension time was 5.60 ± 1.276 (Mean \pm SD) in Nitroglycerine group . Onset time was 13.567 ± 2.176 (Mean \pm SD) in the Isoflurane Group .

107

In our study the time required by NTG to achieve desired SBP was 5.60 ± 1.276 min correlating well with the studies of **Tannieres et al (4-6 min)** and **Shenoy et al (3-6)min.**

Seigle Morach and Green (1986) who could achieve hypotension with Nitroglycerine in 2.1 minutes in most of their cases .

The quick action of Nitroglycerine has been attributed to the direct musculotropic action on the blood vessels , both the venous and arterial systems . (**Goodman and Gillman**) .

In our study the time required by Isoflurane to achive desired SBP is 13.567 ± 2.176 corelating well with studies of **Ankichetty SP , Ponniah M, Thomas S et al (2011) (16 \pm 7 mins)** .

RECOVERY FROM HYPOTENSION : --

Recovery from the hypotensive effects is of special significance as bleeding within the spinal canal cann't be controlled by usual maneuvers like pressure once the skin is closed .

In this series , we found out that the recovery from hypotension was much more faster in the Nitroglycerine group than Isoflurane group . Mean recovery time was 6.33 ± 1.373 in NTG group . It was 13.467 ± 3.098

108

in Isoflurane group group . This difference was found to be significant . ($p < 0.001$) .

Our observations corroborate with the findings of **Yajnik et al (1977)** and **P. Jindal, R. Gupta & J. P. Sharma 2008** who had concluded that onset and recovery from controlled hypotension using organic nitrates is extremely rapid and thus nitrates are definitely superior and more potent hypotensive agents compared to others .

Kadam PP , Saksena SG (1993) and Ankichetty SP , Ponniah M, Thomas S, also had a similar opinion .

DOSE REQUIRED FOR INDUCED HYPOTENSION:-

Both Nitroglycerine and Isoflurane -- based techniques were equally capable of producing controlled hypotension .

The highest concentration used in Isoflurane group to achieve target blood pressure was an inspired concentration of 2.5% - 3.5% and maintenance was achieved with a concentration of 0.5% – 2.5 %. In a study by **Tirelli et al** a concentration of 1 – 2 % of Isoflurane was used in Isoflurane group for the maintenance which is almost similar to our study . In Nitroglycerine group the dose required to maintain the blood pressure at the desired level ranged from average of 0.9482 to 1.716µg/kg/min which

109
is quite similar to the doses used by **Fahmy and Shiraishi et al** . (1.023 ± 0.468).

HEART RATE : --

Table shows the variations in mean heart rate after start of hypotensive agent .

Heart rate was compared in two groups .In group -- N there was a significant increase in heart rate seen from the baseline after induction. There was a highly significant increase in heart rate at 15, 30 minutes after the start of the hypotensive agent and highly significant increase in heart rate almost at a constant level till the end of the surgery when compared with the baseline values.

On comparing intra group we observed that HR had decreased significantly in group - I as compared to group --N . at 15, 30 min.In group - I there was a significant increase in heart rate after induction, at 5min after induction , at the start of hypotensive agent in group I but there was no significant alterations at other time intervals when compared to baseline values.

Kapaln JA , Dunbar RW , 1967 . Kadam PP , Saksena SG , 1993. didn't observe significant change in heart rate when Nitroglycerine was used to control blood pressure during coronary artery surgery . They suggested that gradual reduction in blood pressure by Nitroglycerine prevented a higher

110
increase in heart rate in heart rate . Similar results were reported by **Bale et al , Fahmy NR , 1978 and Dauchet PJ , 1979** . Our study corroborated well with the findings of **P. Jindal, R. Gupta & J. P. Sharma 2008, and Ankichetty SP , Ponniah M, Thomas S, Kumar K et al 2011** that organic nitrates has significantly raised heart rate than inhalational agents.

AMOUNT OF BLOOD LOSS : --

The total amount of blood loss assessed in spine surgeries for Group - N was 219.96 ± 12.992 and for Group - I it was 225.96 ± 11.978 which shows that the mean intraoperative blood loss in Nitroglycerine group was significantly less (p < 0.0259) compared to the Isoflurane group .

These findings conforms to the observations of **Kakiuchi . M (Osaka , Japan 1998)** who demonstrated that the main source of bleeding during posterior spinal surgery is the bone and is venous rather than arterial . The reduced venous pressure decreases intraosseous pressure , drastically reducing bleeding .

111

Nitroglycerine predominantly dilates the venous capacitance vessels causing pooling of blood away from the operating site . The reduced oozing in GTN Group led to a drier and clearer surgical field where as the field was wet and interfered with surgical dissection in the Isoflurane group .

Ankichetty SP , Ponniah M, Thomas S, Kumar K et al 2011 also had a similar opinion .

DURATION OF SURGERY : --

In Group – N the duration of surgery was 134.70 ± 13.754 minutes and in Group – I the duration was 140.83 ± 10.158 minutes .

Though the blood loss is more in Isoflurane group due more oozing, the duration of surgery is statistically not significant . Our observations corroborated well with the study by P. Jindal, R. Gupta & J. P. Sharma (2008) and Ankichetty SP , Ponniah M, Thomas S, Kumar K et al (2011) who had emphasized there is no significant advantage of NTG over Isoflurane inhalational anaesthetic technique in terms of duration of surgery .

112

GRAPH - 7,8,9

The SBP was well maintained in between 80-95mm Hg in groups N and I during the hypotensive period. There was no significant in various groups during the course of hypotension. There was no incidence of rebound hypertension seen in any of the hypotensive group .

There was statistically no significant difference ($p>0.05$) in diastolic blood pressure before induction in all the groups. Average 33-40% reduction in DBP was seen from the baseline values in all the hypotensive groups. When compared to baseline, the DBP at 15, 30, 45,60min till the drug discontinuation the variation was highly significant due to deliberate hypotension.

Mean arterial pressure (MAP) decreased from 31-35% in all the two groups. There was no significant difference in MAP in two groups during the course of hypotension.

COMPLICATIONS : --

The complications that had or might have been observed were shown in the table .

113

Prolonged drowsiness for up to 2-4 hrs was seen in cases in 2 cases in Isoflurane group compared to nil in group – N .

None of the patients in two groups had postoperative nausea and vomiting . This could be due to prophylactic administration of ondansetron in both the groups .

There were no signs of cerebral dysfunction in either of the groups . Not a single case of allergic manifestations was observed in this series . Rebound hypotension was also not observed in any patients after discontinuation of

respective hypotensive agent . There was no evidence of tolerance or tachyphylaxis to the action of Nitroglycerine .
No other life threatening complications was observed .

Summary

60 cases from Neurosurgery Inpatients resister of S.C.B. Medical college and Hospital , Cuttack , posted for elective spine surgery (Lumber laminectomy and decompression ; posterior approach) where controlled hypotension was indicated were taken up for this study . The cases were divided into two demographically identical groups. In one group hypotension was induced by using Nitroglycerine infusion in the dose range 1 – 10 µg /kg / min and in other group , Isoflurane inhalation was used in flutech Vaporiser in a concentration of 1 – 4 % . All the patients were administered general . SBP was lowered to 80 – 100 mmHg before skin incision and maintained throughout the surgery by titrating the doses of drugs in both the groups .

The following points were noted for each drug and tabulated .

1. Speed of onset of hypotension .
2. Intraoperative heart rate variations .
3. Speed of recovery from hypotension .
4. Mean blood loss .
5. Duration of surgery .
6. No of patients requiring blood transfusion .

116

7. Surgeon's scoring .
8. Post operative complications

Observations were compared between the two groups using standard statistical analysis .

The following conclusions were derived from the study .

- Nitroglycerine induces hypotension more rapidly and speed of recovery from hypotension is also faster .
- Intra- operative blood loss is significantly less in Nitroglycerine group as oozing is more in isoflurane group than Nitroglycerine group .
- Because of the reduced blood loss the need for blood transfusion is less in Nitroglycerine group .
- In both the groups all the patients belong to Grade 3 and below which denotes highly acceptable surgical field as far as the surgeon concerned , However the no. of patients in Grade 3 are more in Isoflurane group than Nitroglycerine group.

Conclusiuon

Though controlled hypotension is produced effectively and surgical field is acceptable with isoflurane, Nitroglycerine is a safe and potent drug to achieve controlled hypotension during surgical operations. □ □

Bibliography

1. Adams et al;Techniques of vascular control for deliberate hypotension during anaesthesia. BJA;1975;47;777-792.
2. Aitken D, West D, Smith F, et al: Cyanide toxicity following nitroprusside induced hypotension. Can Anaesth Soc J 24:651, 1977
3. Anderson et al;Deliberate hypotensive anaesthesia for orthognathic surgery:controlled pharmacologic manipulation of cardiovascular physiology. Int J Adult Orthodon Orthognath Surg;1986;1;133-159.
4. Ankichetty SP, Ponniah M, Cherian yr. Thomas S. Kumar K, et al. Comparison ot totalIntravenous anesthesia using propofol and Inl alational anesthesia using isoflurane forcontrolled hypotension In functional endoscopic sinus surgery_ J Anaesth' (Clin Pharmacol 2011 ;27:328-32.)
5. Artru AA, Nugent M, Michenfelder JD: Enflurane causes a prolonged and reversible increase in the rate of CSF production in the dog. Anesthesiology 57:255, 1982.
6. Beaussier M,Paugam C,Deriaz H.Haemodynamic stability during moderate hypotensive anaesthesia for spinal surgery.A comparison between desflurane and isoflurane. Acta Anaesthesiol Scand 2000;44: 1154-59. (s)
7. Beierholm.W.E, Bregaard Sorenson. M, Sroczyński. Z et al. - haemodynamic changes during sodium nitroprusside induced hypotension and halothane/nitous oxide anaesthesia. acta anaesthesiology scandinavia 1983 april; 27(2): 99-1031.
8. Bergman.S, Hoffman W.E, Gans B.J, Miletich D.J et al. blood flow to oral tissues; and experimental study with enflurane, sodium nitroprusside and nitroglycerin. journal of maxillo facial surgery.1982 jan; 40 (1): 13
9. Boezaart AP, van der Merwe J, Coetzee A. Comparison of sodium nitroprusside- and esmolol-induced controlled hypotension for functional endoscopic sinus surgery. can J Anaesth 1995;42:373-6.
10. Briggs JB, Cook HW. The value of an accurate knowledge of arterial blood pressure to the clinician. Md Med J. 1903; 46:1-13
11. Bromage PR. Vascular hyotension in 107 cases of epidural analgesia. Anaesthesia 1951; 6:26. (s) 119
12. Buffington CW, Romson JL, Levine A, et al: Isoflurane induces coronary steal in a canine model of chronic coronary occlusion. Anesthesiology 66:280, 1987.
13. Campkin T.V. Flinn R.M.Isoflurane: its use to induce hypotension in

neurosurgical patients. *European Journal of Anaesthesiology* 1986 sep; 3(5): 39-40

14. Casthely PA, Lear S, Cottrell JE, et al: Intrapulmonary shunting during induced hypotension. *Anesth Analg* 61:231, 1982.

15. Chan.W, Smith D.E, effects of hypotensive anaesthesia in anterior maxillary osteotomy. *Journal of Oral Surgery* 1980 jul; 38(7): 504-86.

16. Chen RYZ, Matteo RS, Fan F-C, et al: Resetting of baroreflex sensitivity after induced hypotension. *Anesthesiology* 56:29, 1982.

17. Cnlund M.G, Ahistedt B.I, Anderson I.G et al. induced hypotension may influence blood loss in orthognathic surgery, but it is not crucial. *Scand J Plast Reconstr Surg Hand Surg*.1997 dec: 31(4): 311-79.

18. Crystal GJ, Salem MR: Myocardial and systemic hemodynamics during isovolemic hemodilution alone and combined with nitroprusside-induced controlled hypotension. *Anesth Analg* 72:227, 1991.

19. Cushing H. On routine determinations of arterial tension in operating room and clinic. *Boston Medical and Surgical Journal*. 1903; 148:250-2.

20. Cushing H: Tumors of the Nervus Acusticus. Philadelphia, WB Saunders,1917.

21. Didier EP, Clagett OT, Theye RA: Cardiac performance during controlled hypotension. *Anesth Analg* 44:379, 1965.

22. Donald JR: Induced hypotension and blood loss during surgery. *J R Soc Med* 75:149, 1982.

23. Eckenhoff JE, Crompton JR, Larson A, et al: Assessment of the cerebral effects of deliberate hypotension by psychological measurements. *Lancet* 2:711, 1969.

24. Eckenhoff JE, Rich JC: Clinical experiences with deliberate hypotension. *Anesth Analg* 45:21, 1966. *Anesthesiology* 48:87, 1978.

25. Ekback G, Aexlsson K, Ryttberg I et al. tranexamic acid reduces loss in total hip replacement surgery. *Anesthesia Analgesia* 2000 nov; 91(5): 1124- 308.

26. Enderby GEH. Controlled circulation with hypotensive drugs and posture to reduce bleeding during surgery. Preliminary results with pentamethonium iodide. *Lancet* 1950;1:1145. (s).

27. Fahmy et al ;Indications and contraindications for deliberate hypotension.;*Int Anesthsiology Clin*;1979;17;175-187.

28. Fahmy N.R. nitroglycerin as a hypotensive drug during general anaesthesia. *anaesthesiology* 1978 jul; 49(1): 17-2010.

29. Fahmy NR: Nitroglycerin as a hypotensive drug during general anesthesia. *Anesthesiology* 49:17, 1978.

120

30. Fromme et al; Controlled hypotension for orthognathic surgery. *Anesth Analg*;1986;65;683-686.

31. Fulton JF. Harvey Cushing: A Biography. Springfield, IL: Charles C. Thomas; 1946:178-9, 184.

32. Gallagher, Milliken. induced hypotension for orthognathic surgery. *Journal of Oral Surg* 1979 jan; 37(1): 47-5112.

33. Ganong WF: Review of Medical Physiology, 14th ed. E. Norwalk, Appleton & Lange, 1989, p 600.
34. Gardner JW. The control of bleeding during operation by induced hypotension .JAMA1946;132: 572. (s)
35. Goldberg ME, McNulty SE, Azad SS. A comparison of labetolol and nitroprusside for inducing hypotension during major surgery. Anaesthesia and Analgesia 1990;70: 537-42. (s)
36. Golia J.K, Woo R, Farole a et al. nitroglycerin- controlled circulation in orthognathic surgery. journal of oral maxillofacial surgery 1985 may; 43(5): 342-513.
37. Goodman and gillman. the pharmacology- basics of therapeutics.
38. Goodman EH, Howell AA. Clinical studies in the auscultatory method of determining blood-pressure. University of Pennsylvania Bulletin. 1910-11; 23:469-75
39. Goran zellin, Dars Rasmusson, Jan Palsson et al. evaluation of hemorrhage depressors on blood loss during orthognathic surgery a retrospective study journal of oral maxillofacial surgery 2004; 62:662-666.15.
40. Gourdeau M, Martin R, Lamarche Y, Tetreault I; oscillometry and direct blood pressure: a comparative clinical study during deliberate hypotension. canadian anaesthetist society. 1986 may; 33(3pt1): 300-7
41. Greene NM. Hypotensive spinal anaesthesia. Surg Gynecol Obstet 1952; 95: 331. (s)
42. Griffiths HWC, Gillies J: Thoraco-lumbar splanchnicectomy and sympathectomy: Anaesthetic procedure. Anaesthesia 3:134, 1948.1
43. Guggiari M, Dageouf, Lienhart A et al. use of nitroglycerine to produce controlled decreases in mean arterial pressure to less than 50 mm hg. british journal of anaesthesia 1985 feb; 57(2): 142-718.
44. Guyuron B, Vaughan C, Schlecter B. the role of ddavp (desmopressin) in orthognathic surgery. anesth analg: 2000 nov; 91(5): 1124-3019.
45. Hoffman W.E, Bergman S, Miletich D J et al. regional vascular changes during hypotensive anaesthesia. journal of cardio vasc pharmacol.1982 march- april; 4(2): 31020.
46. Jacobi K.E, Bohn B.E, Riickaver A.J et al. moderate controlled hypotension with sodium nitroprusside does not improve surgical 121 conditions or decrease blood loss in endoscopic sinus surgery. journal of clinical anaesthesia 2000 may; 12(3): 202-721.
47. Jam AM, Gelb AW. cardiovascular effects of isoflurane – induced hypotension for cerebral aneurysm surgery. anesth analg. 1993 aug; 62 (8): 742-826.
48. Jean-Marc Bernard, Micheal Pinaud, Thierry Francois et al. deliberate hypotension with nicardipine or nitroprusside during hip arthroplasty, 1991; 73:341-51
49. John E. Moenning, David A. Bussard, Thomas H.Lapp et al. average blood transfusion in 506 orthognathic surgical procedures. journal of oral

- and maxillofacial surgery 1995; 53:880-883,24.
50. Kadam PP, Saksena SG, Jagtap SR, Pantavaidya SM. Hypotensive anaesthesia for spine surgery -nitroglycerine versus halothane .J Postgrad Med 1993; 39:26-28. (s)
51. Kaplan JA: The role of nicardipine during anesthesia and surgery. Clin Ther 11:84, 1985
52. Kerr A.R. anaesthesia with profound hypotension for middle ear surgery. british journal of oral and maxillofacial surgery 1977 may; 49(5): 447-5225.
53. Khambatta HJ, Stone JG, Khan E: Propranolol alters renin release during nitroprusside-induced hypotension and prevents hypertension on discontinuation of nitroprusside. Anesth Analg 60:569, 1981.
54. Korotkoff NC. On methods of studying blood pressure. Izvestiia Voennomedistinskite Akademiia. 1905; 11:365
55. Larsen et al ;Cerebral blood flow and metabolism during isoflurane induced hypotension in patients subjected to surgery for cerebral aneurysms.;BJA;1987;59;1204-1207.
56. Larson AG: Deliberate hypotension. Anesthesiology 25:682, 1964.
57. Lassen NA, Christensen MS: Physiology of cerebral blood flow. Br J Anaesth 48:719, 1976.
58. Leigh J M. the history of controlled hypotension. british journal of anaesthesia 1975 july; 47(7): 745- 74927.
59. Lessard et al ;Isoflurane induced hypotension in orthognathic surgery.;AnesthAnalg;1989;69;379-383.
60. Little DM Jr. Induced hypotensionduring anaesthesia and surgery .Anaesthesiology 1955;16:320-332. (s)
61. Mac Rae W.R. induced hypotension. BR J Hosp med 1985 Jun; 33(6): 341-328.
62. Mandai P. Isoflurane anesthesia for functional endoscopic sinus surgery. Indian J Anaesth 2003;47:37-40.
122
63. Mandel RJ, Brown MD,Mc Collough NC,Pallares V,Varlotta R.Hypotensive anaesthesia and autotransfusion in spinal surgery. Cl. Orthop Rel Resear 1981;154:27-33. (s)
64. Martin R. lessard, Claude a et al. isoflurane- induced hypotension in orthognathic surgery. anesth analg.1989; 69:379-8329.
65. Michael.E.Goldberg,.Mcnulty S.E et al. a comparison of labetalol and nitroprusside for inducing hypotension during major surgery 1990;vol: 70:537-42
66. Michenfelder JD, Theye RA: Canine systemic and cerebral effects of hypotension induced by hemorrhage, trimetaphan, halothane, or nitroprusside. Anesthesiology 46:188, 1977.
67. Murat I,Delleur MM,Loose JP,Saint Maurice C. Controlled hypotension during posterior vertebral arthrodesis ; value of an isoflurane - nitroglycerine combinaiton.Anaesth Analg 1981; 60:513-16. (s)
68. Murtagh GP.Controlled hypotension with halothane .Anaesthesia

1960;15:235. (s)

69. Myron Yaster, Raymonds. Simmons, Vernon t. Tolo et al. a comparison of nitroglycerin and nitroprusside for inducing hypotension in children. *anesthesiology* 1986; 65:175-179,31.

70. Nabil samman, LIM Kwong Cheung, Antonio C.K.Tong et al. blood loss and transfusion requirements in orthognathic surgery. *journal of oral and maxillofacial surgery* 1996; 54:21-24,32.

71. Nair S, Collins M, Hung P, Rees G, Close D, Wormald PJ. The effect of beta-blocker premedication on the surgical field during endoscopic sinus surgery. *Laryngoscope* 2004;114:1042-6.

72. Nakazawa K,Taneyama C,Benson KT,Unruh GK, Goto H. Mixtures of sodium nitroprusside and trimethaphan for the induction of hypotension. *Anaesthesia and analgesia*1991; 73:59-63. (s)

73. Neil M.C, wald D.E, Bennet E.J et al. controlled hypotensive anesthesia in scoliosis surgery. *bone joint surgery am* 1974 sep; 56(6): 1167-7233.

74. Ornstein E,Young WL,Ostapkovich N,Matteo RS, Diaz J. Deliberate hypotension in patients with intracranial arteriovenous malformations:esmolol compared with isoflurane and sodium nitroprusside. *Anaesthesia and Analgesia* 1991;72:639-44. (s)

75. P. Jindal, R. Gupta & J. P. Sharma : Is a combination of Isoflurane with nitroglycerine better than halothane with nitroglycerine for controlled hypotension in spine surgery: A comparative clinical evaluation? . *The Internet Journal of Anesthesiology*.

76. Pasch T, HUK W. Cerebral complications following induced hypotension. *Eur J anaesthesiol* 1986 Jul; 3(4): 299-31234.

123

77. Patel H: Experience with the cerebral function monitor during deliberate hypotension. *Br J Anaesth* 53:639, 1981.

78. Petrozza et al ;Induced hypotension.; *Int Anesthesiology Clin*;1990;28;223-229.

79. Praveen K, Narayanan V, Muthusekhar M.R et al. hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study. *british journal of oral and maxillofacial surgery* 2001; 138-14036.

80. Precious D.S, Splinter W, Bosco D. induced hypotensive anesthesia for adolescent orthognathic surgery patients. *journal of oral and maxillofacial surgery* 1996; 54:683-684,

81. Price ML, Walmsley A, Swaine C, Ponte J. Comparison of total intravenous anaesthetic technique using a propofol infusion, with an inhalational technique using enflurane for day case surgery. *Anaesthesia* 1998;43:84-7.

82. Priebe H-J: Isoflurane causes more severe regional myocardial dysfunction than halothane in dogs with critical coronary artery stenosis. *Anesthesiology* 69:72, 1988.

83. Prys-Roberts C, Lloyd JW, Fisher A, et al: Deliberate profound hypotension induced with halothane: Studies of haemodynamics and pulmonary gas exchange. *Br J Anaesth* 46:105, 1974.

84. Ramsey M. blood pressure monitoring; automated oscillometric devices. *journal of clinical monitoring* 1991 jan; 7(1): 56-6738.
85. Ray M, Bhattacharjee Dp, Hajra B, Pal R, Chatterjee N. Effect of clonidine and magnesium sulphate on anaesthetic consumption, haemodynamics and postoperative recovery: A comparative study. *Indian J Anaesth* 2010;54:137-41.
86. Reiz S, Balfors E, Sorensen MB, et al: Isoflurane—a powerful coronary vasodilator in patients with coronary artery disease. *Anesthesiology* 59:91, 1983.
87. Riva-Rocci S. Un nuovo sfigmomanometro. *Gazzetta Medica di Torino*. 1896; 47:981-96, 1001-17
88. Robert M. Dolman, Kenneth C. Bentle et al. The effect of hypotensive anesthesia on blood loss and operative time during lefort i osteotomies. *journal of oral and maxillofacial surgery* 2000; 58:840,39.
89. Rodrigo et al ;In Rodrigo et al ;duced hypotension during anaesthesia with special reference to orthognathic surgery.;*Anesth Prog*;1995;42;41-58.
90. Rohling R.G, Zimmerman A.P, Biro P et al. Alternative methods for reduction of blood loss during elective orthognathic surgery. *international journal of adult orthodontics and orthognathic surgery* 1999; 14:77-8240. 124
91. Rout A. Anaesthesia for Neurosugery .In: Ramamurthy B,Tandon PN, Ramamurthy R,Sridhar K.Textbook of Neurosurgery .2nd ed.B I Churchill Livingstone 1996:1346-1357. (s)
92. Ryu JH, Sohn IS, Do SH. Controlled hypotension for middle ear surgery: A comparison between remifentanil and magnesium sulphate. *Br J Anaesth* 2009;229: 1-6.
93. Schaberg S.J, Kelly J.R, Terry b.G et al. Oral-Facial corrective surgery. *journal of oral surgery* 1976 feb; 34(2): 147-5642.
94. Shenoy V,Rajan R.K., Rao G,Bhatt G,Upadhya M. Deliberate hypotension-a comparative study using nitroglycerine and sodium nitroprusside .*Ind J Anaesth* 2000;44:38-9. (s)
95. Shiraishi Y,Mochizuki T,Ikeda K.Oxygen uptake and carbon dioxide elimination during controlled hypotension induced by prostaglandin E1 or nitroglycerine.*Br. J. Anaesth* 1994; 72:439-442. (s)
96. Simpson DA, Ireland J: Hypotensive and normotensive anaesthesia in total hip replacement. A comparative study. *Br J Clin Pract* 37:16, 1983.
97. Sivarajan M, Amory DW, Everett GB, et al: Blood pressure, not cardiac output, determines blood loss during induced hypotension. *Anesth Analg* 59:203, 1980.
98. Schindler I, et al Nitroglycerine and isoflurenane for controlled hypotension in maxilofacial surgery. *Acta Anaesthesiology* 1994.
99. Sollevi a.hypotensive anaesthesia and blood loss. *acta anaesthesiol scand supply* 1988; 89:39-43
100. Stephen E mc Nulty.induced hypotension during head and neck surgery. *anaesthesiology clinics of north america*, 1993 sep; 11(3)45.
101. Stewart. A, Newman. L, Sneddon. K, Harris. M.Aprootinin reduces blood

- loss and the need for transtusion in orthognathic surgery. british journal of oral & maxillofacial surgery 2001; 30:365-370.46.
102. Slack WK et al;Organ perfusion during controlled hypotension. al;BJA;1975;47;793-798.
103. Suwa K, Hedley-Whyte J, Bendixen HH: Circulation and physiologic dead space changes on controlling the ventilation of dogs. J Appl Physiol 21:1855, 1966.
104. Tannieres ML,Chelly J, Dermone P,Visto A,Vourch P. Clinical use of nitroglycerine as ahypotensive agent during general anaesthesia. Anesth Anlag 1979;36:139-145. (s)
105. Thompson GE, Miller RD, Stevens WC, et al: Hypotensive anesthesia for total hip arthroplasty: A study of blood loss and organ function (brain, heart, liver and kidney). Anesthesiology 48:91, 1978.
106. Thulin T, Andersson G, Schersten B. Measurement of blood-pressure-a routine test in need of standardization. Postgrad Med J. 1975; 51:390-5. 125
107. Tirelli G, Bigarini S, Russolo M, Lucangelo U, Gullo A. Total intravenous anaesthesia in endoscopic sinus-nasal surgery. Acta Otorhinolaryngol Ital 2004;24:137-44.
108. Tremper K.K, Waxman K.S, Applebaum R.A et al. Transcutaneous PO2 monitoring during sodium nitroprusside infusion. crit care med.1985 feb; 13(2): 65-748.
109. Van DE Perre JP, Stoelinga P.J, Blijdorp P.A et al. perioperative morbidity in maxillofacial surgery: a retrospective study. j craniomaxillofac surg1996 oct; 24(5): 263-7049.
110. Vandesteene, Mouawad, Noterman et al. use of sodium nitroprusside in neurosurgical cases during anesthesia with enflurane. acta anaesthesiology 1980; 31:suppl: 85-92
111. Wad et al;Deliberate hypotension during head and neck surgery. al;Head and neck surgery;1980;2;185-195.
112. Ward C.F. , Alfery David D, Saidman L. J., Waldman J. Deliberate hypotension in head and neck surgery. Wiley Intersciences 2006; 3:185-195. (s)
113. Washburn MC,Hyer RL. Deliberate hypotension for elective major maxillofacial surgery:a balance halothane and morphine technique.J Maxillofac Surg 1982; 10(1): 50-55. (s)
114. Wildsmith JAW, Drummond GB, MacRae WR: Blood-gas changes during induced hypotension with sodium nitroprusside. Br J Anaesth 47:1205, 1975.
115. Yu c.n.f, chow t.k, kwan a.s.k. et al intra-operative blood loss and operating time in orthognathic surgery using induced hypotensive general anaesthesia: prospective study. hong kong medical journal 2000; 6:307-11
116. Wyman et al , Induced hypotension using Hexamethonium and pentamethidium. Anaesthesiology 60,541, 1953.
117. Zall S,Milocco I,Ricksten S.E. Effects of adenosine on myocardial blood

flow and metabolism after coronary artery bypass surgery. Anaesthesia and Analgesia 1991;73:689-95. (s).

□□□