

Original research article

**A STUDY OF ASSOCIATION OF *H. PYLORI* WITH
GALLSTONE DISEASE**

¹G Sai Anurag, ²Dr. Harischandra B, ³Dr. Rekha PD, ⁴Dr. Vajrang Akula

^{1,4}Senior Resident, Department of General Surgery, Yenepoya Medical College,
Karnataka, India

²Professor and HOD, Department of General Surgery, Yenepoya Medical College,
Karnataka, India

³Professor and Director, Yenepoya Medical College, Karnataka, India

Corresponding Author:

Dr. Vajrang Akula

Abstract

The prevalence of gall bladder stones in our country varies between 10-20% of the population, with approximately 4.3% of individuals experiencing symptoms. The presence of an imbalance in the composition of bile, bile stasis, or gallbladder infections might potentially result in the development of gallstones. Nevertheless, numerous research have examined and documented the association between *H. pylori* and disorders affecting organs outside the stomach and duodenum.

Keywords: Gall bladder, *Helicobacter pylori*, association

Introduction

The gallbladder is an oblong organ located on the inferior surface of the liver in the gallbladder fossa. Its primary function is to store bile ^[1]. Imbalances in the composition of bile can result in the development of gallstones ^[2]. The gallbladder is susceptible to the development of various conditions such as cancer and inflammation ^[3]. The prevalence of gall bladder stones in our country varies between 10-20% of the population, with approximately 4.3% of individuals experiencing symptoms. Gallstones can occur when there is an imbalance in the composition of bile, stasis of bile, or infections in the gallbladder ^[4, 5]. The gallbladder is susceptible to several disorders such as cancer and inflammation ^[7, 8]. The prevalence of gallstone disease in India is relatively high, ranging from 10-20% ^[9]. Prevalence is higher among females. Recently, many scientists have become interested in studying the relationship between *H. pylori* infection and different hepatobiliary diseases. However, due to the intricate pathophysiologic mechanisms involved in this infection, the diagnostic procedures for detecting *H. pylori* in extra gastric specimens have not met the desired standards of accuracy ^[11, 23]. Nevertheless, the association between *H. pylori* and disorders affecting organs outside the stomach and duodenum has been extensively examined and documented in several studies ^[24, 26].

Aim and objective

To find out the incidence of *H. pylori* infection in the gallbladder of patients with symptomatic gallstone disease.

Materials and Methods

Study design: observational study

Study setting: Yenepoya Medical College Hospital (YMCH) is a 900 bedded tertiary care teaching hospital situated in Deralakatte, a suburban locality of Mangaluru, Dakshina Kannada. It provides general and specialist healthcare to the coastal and central parts of Karnataka and northern part of Kerala.

Type of study: Prospective study.

Study Population

Patients undergoing cholecystectomy for symptomatic gallstone disease in Yenepoya Medical College Hospital.

Study period

This was conducted between May 2021 and October 2022.

Inclusion criteria

- Patients admitted with symptomatic gallstone disease undergoing cholecystectomy at Yenepoya medical college hospital.
- Age of a patients more than 18 years.

Exclusion criteria

- Patients who have undergone any previous hepatobiliary surgery or intervention.
- Patients who received *H. pylori* eradication treatment in the last 1 year.
- Intervention.

No intervention other than the surgery needed. Sampling procedure: Convenient sampling.

Randomization: nil.

Purposive sampling: Nil

Sample Size: 42.

Sample Size formula: $n = Z^2 \cdot \frac{p \cdot q}{d^2}$ at 5% level of significance and 54.2% of proportion (from related article) with 15% of margin of error, the total sample size is 42.

- Study was conducted in YMHC from May 2021 - August 2022.
- After the ethical committee and SRB clearance, the study was started.
- Patients were selected according to the criteria; informed consent was taken from every patient.

Method of collection of data

- All patients who meet the predefined criteria were counseled regarding the study following which those patients who give a written informed consent on their free will without any monetary gain will be chosen for the study. They then underwent complete medical history and clinical evaluation a pre anesthetic checkup. After being cleared from the anesthetists, they underwent surgery.
- Each patient underwent an endoscopy to rule out any stomach or duodenal pathology as the cause for upper abdominal pain. A ultrasound was done to look at the status of the gallbladder. On the day planned for surgery, the patient is kept nil by mouth for at least 8 hours.

Following extraction of the gall bladder the following samples are collected

Gallbladder mucosa biopsy, for Rapid urease testing, where the principle that *H. pylori* ferment urea is utilized to identify the presence of *H. pylori*. This test was done using a RUT dry test kit, A small 2-3 mm size of mucosal tissue is taken and introduced onto the yellow coloured media on the kit. If there is a change in the colour of the media from yellow to pink then it is called as a positive test result.

Statistical Analysis

- Descriptive Statistics.
- Mean and Standard deviation for categorical data Frequency and percentage for categorical data.
- Pearson's correlation was used to find out correlation between RUT and DNA.

Results


This Prospective study observational study was conducted at Yenepoya Medical College Hospital (YMCH) is a 900 bedded tertiary care teaching hospital situated in Deralakatte, a suburban locality of Mangaluru, Dakshina Kannada with patient population from local, neighboring districts as well as northern part of Kerala.

The study period was between May 2021 and October 2022 on Patients undergoing cholecystectomy for symptomatic cholelithiasis in Yenepoya Medical College Hospital.

Age

Table 1: Age

Age in years	Case number	Percent of cases
21-30 YEARS	08	19.09%
31-40 YEARS	14	33.33%
41-50 YEARS	14	33.33%
51-60 YEARS	03	07.14%
61-70 YEARS	03	07.14%
TOTAL	42	100.00%




Graph 1: Age

The age of the cases in the study ranged between 22 years and 67 years, the mean age of the cases was 41.28 years SD +/- 10.87 years.

Gender

Table 2: Gender

Gender	Case number	Percent of cases
Females	33	78.57%
Males	09	21.42%
Total	42	100.00%




Graph 2: Gender

33 (78.57%) cases were females and 09 (21.42%) cases were males. Chi square p=0.023 significant, male to female ratio was 0.27:1

***H. Pylori* by rapid urease test**

Table 3: *H. Pylori* by rapid urease test

Rapid urease test	Case number	Percent of cases
Positive for <i>H. Pylori</i>	24	58.54%
Negative for <i>H. Pylori</i>	18	43.90%



Graph 3: *H. Pylori* by rapid urease test

Rapid urease test results showed that 58.54% (24) of the gallbladder mucosal specimens were positive for *H. pylori* and 43.9% (18) of the gallbladder mucosal

specimens were negative for presence of *H. pylori*.

Discussion

This prospective observational study was carried out at Yenepoya Medical College Hospital (YMCH) in Deralakatte, Mangaluru, Dakshina Karnataka, from May 2021 to October 2022. The study focused on patients receiving cholecystectomy for symptomatic cholelithiasis. The following is a summary of the results. The age spanned from 22 to 67 years, with a mean age of 41 years and a standard deviation of 10.87 years. 31 instances, accounting for 73.81% of the total, were females. Out of the total number of cases, 11 cases, which accounts for 26.19% of the total, were men. The chi-square test yielded a p-value of 0.023, indicating statistical significance. The male to female ratio was 0.33:1.

H. pylori the quick urease detection method had a sensitivity of 80% and a specificity of 60%. The disease prevalence was 88.24%. The positive predictive value was 92.31%, while the negative predictive value was 35%. The accuracy of the test was 76.47%. The prevalence of gallstones rises with age. The prevalence of this condition is higher in women than in men, with a ratio of 4:1. Additionally, around 50% of patients do not exhibit any symptoms. The development of gallstones is influenced by multiple factors. Alongside alterations in bile composition or bile stasis, the primary cause of gallstone production appears to be the presence of an infectious agent. The phrase coined by Moynihan, which states that a gallstone is a memorial created within the organism, remains accurate in the present time. Laparoscopic cholecystectomy (LC) is a widely performed surgical operation that is considered the most effective treatment for symptomatic gallstones.⁸ Gallstone disease is prevalent in middle-aged females, with its occurrence rising between the ages of 30 to 50. The prevalence of gall bladder stones in our country varies between 10% and 20% of the population, and it causes symptoms in approximately 4.3% of individuals.

Conclusion

In this study it was found that there was presence of *H. pylori* in the gallbladder mucosa of patients with cholelithiasis. Rapid urease testing showed presence of *H. pylori* in 24 patients (54.54%).

References

1. Frierson Jr HF. The gross anatomy and histology of the gallbladder, extrahepatic bile ducts, Vaterian system, and minor papilla. *The American journal of surgical pathology*. 1989 Feb 1;13(2):146-162.
2. Toouli J, Bhandari M. Anatomy and physiology of the biliary tree and gallbladder. *Diseases of the Gallbladder and Bile Ducts Diagnosis and Treatment*, Second Edition. Massachusetts: Blackwell Publishing; c2006. p. 3-21.
3. Song ST, Shi J, Wang XH, Guo YB, Hu PF, Zhu F, *et al*. Prevalence and risk factors for gallstone disease: A population-based cross-sectional study. *Journal of digestive diseases*. 2020 Apr;21(4):237-245.
4. Pogorelic Z, Aralica M, Jukic M, Zitko V, Despot R, Juric I, *et al*. Gallbladder disease in children: a 20-year single-center experience. *Indian pediatrics*. 2019

May;56(5):384-386.

5. Çerçi SS, Özbek FM, Çerçi C, Baykal B, Eroğlu HE, Baykal Z, *et al.* Gallbladder function and dynamics of bile flow in asymptomatic gallstone disease. *World journal of gastroenterology: WJG.* 2009 Jun 14;15(22):2763.
6. Toouli J, Bhandari M. Anatomy and physiology of the biliary tree and gallbladder. *Diseases of the Gallbladder and Bile Ducts Diagnosis and Treatment, Second Edition.* Massachusetts: Blackwell Publishing; c2006. p. 3-21.
7. Song ST, Shi J, Wang XH, Guo YB, Hu PF, Zhu F, *et al.* Prevalence and risk factors for gallstone disease: A population-based cross-sectional study. *Journal of digestive diseases.* 2020 Apr;21(4):237-245.
8. Pogorelic Z, Aralica M, Jukic M, Zitko V, Despot R, Juric I, *et al.* Gallbladder disease in children: A 20-year single-center experience. *Indian pediatrics.* 2019 May;56(5):384-386.
9. Stinton LM, Shaffer EA. Epidemiology of gallbladder disease: cholelithiasis and cancer. *Gut and liver.* 2012 Apr;6(2):172-173.
10. Khuroo MS, Mahajan R, Zargar SA, Javid G, Sapru S. Prevalence of biliary tract disease in India: A sonographic study in adult population in Kashmir. *Gut.* 1989 Feb 1;30(2):201-205
11. Cesarani F, Martina MC, Boano R, Grilletto R, D'Amicone E, Venturi C, *et al.* Multidetector CT study of gallbladder stones in a wrapped Egyptian mummy. *RadioGraphics.* 2009 Jul;29(4):1191-1194.
12. Greydanus DE, Joav Merrick MD. Liver and gallbladder: A historical perspective. *International Journal of Child Health and Human Development.* 2020 Jul 1;13(3):229-261.
13. Cuschieri A, Buess G. Introduction and historical aspects. In *Operative manual of endoscopic surgery* Springer, Berlin, Heidelberg; c1992. p. 1-5.
14. Gembal P, Milik K, Ździebło J, Kęsik J, Zubilewicz T. The history of open and laparoscopic cholecystectomy. *Polish Surgery/Chirurgia Polska.* 2007 Jun 1;9(2):1-9.
15. De U. Evolution of cholecystectomy: A tribute to Carl August Langenbuch. *Indian J Surg.* 2004 Mar 1;66(2):97-100.
16. Chalkoo M, Ahanger S. The Historical Perspective, Current Advancements and Innovations in Laparoscopic Cholecystectomy. *J Pak Med Stud.* 2012 Sep 1;2(3):5-6.
17. Adkins RB, Chapman WC, Reddy VS. Embryology, anatomy, and surgical applications of the extrahepatic biliary system. *Surgical Clinics.* 2000 Feb 1;80(1):363-379.
18. Davenport M. The Liver and Gallbladder. In *Clinical Embryology.* Springer, Cham. 2019:353-363.
19. Dachman AH, Schneck C. Embryology of the gallbladder. In *Imaging Atlas.* CRC Press; c2018 Jan 18. p. 13-20.
20. Saldinger PF, Bellorin-Marín OE. Anatomy, Embryology, Anomalies, and Physiology of the Biliary Tract. In *Shackelford's Surgery of the Alimentary Tract.* 2019 Jan;2:1249-1266.
21. Hull NC, Schooler GR, Lee EY. Bile Duct and Gallbladder. In *Pediatric Body MRI.*

- Springer, Cham; c2020. p. 235-253.
22. Rao BN. Analysis of Gallbladder Diseases Diagnosed at a Tertiary Care Hospital: A Retrospective Study. *International Journal of Contemporary Medicine*. 2017 Jan;5(1):156.
 23. Van Gulik TM, Van Den Esschert JW. James Cantlie's early messages for hepatic surgeons: how the concept of pre-operative portal vein occlusion was defined; c2010. p. 81-83.
 24. Khan KS, Sajid MA, McMahan RK, Mahmud S, Nassar AH. Hartmann's Pouch Stones and Laparoscopic Cholecystectomy: The Challenges and the Solutions. *JLS: Journal of the Society of Laparoscopic & Robotic Surgeons*, 2020 Jul, 24(3).
 25. Bergamaschi R, Ignjatovic D. More than two structures in Calot's triangle. *Surgical endoscopy*. 2000 Apr;14(4):354-357.
 26. Abdalla S, Pierre S, Ellis H. Calot's triangle. *Clinical anatomy*. 2013 May;26(4):493-501.