

ORIGINAL RESEARCH

Perioperative consideration in management of severely bleeding coagulopathic patient**¹Sulaiman Hussain Bakr, ²Basman Ali Abbas Hamandi, ³Ahmed Farrag Ali Khaleel Al Alwi**¹Specialist Anesthesia, Burjeel Day Surgery Centre - Al Reem, Abudhabi²Anesthesia Specialist, FIRST IVF Fertility Center, Abudhabi³Specialist Anesthetist, Healthpoint Hospital / Mubadala Health / M42, Abudhabi**Corresponding Author**

Sulaiman Hussain Bakr

Specialist Anesthesia, Burjeel Day Surgery Centre - Al Reem, Abudhabi

Email: sssbakr@gmail.com

Received: 23 November, 2023

Accepted: 26 December, 2023

Abstract**Background:** Perioperative bleeding in severely coagulopathic patients poses significant challenges in clinical management and impacts patient outcomes. Tailored perioperative strategies are essential for optimizing care in this complex cohort.**Objective:** This clinical study aimed to investigate the correlation between preoperative coagulation profiles, surgical interventions, and bleeding outcomes in severely bleeding coagulopathic patients within a tertiary care center.**Methods:** A retrospective analysis was conducted on 73 severely bleeding coagulopathic patients admitted to the center between 2020-2023. Data collection included demographic characteristics, preoperative coagulation profiles, details of surgical interventions, and postoperative outcomes. Statistical analyses were performed to assess the relationship between preoperative parameters, surgical procedures, and bleeding outcomes.**Results:** Preoperative coagulation profiles, including prolonged PT, aPTT, elevated INR, and decreased platelet counts, correlated significantly with the severity of perioperative bleeding. Different surgical procedures exhibited varying extents of bleeding, with orthopedic and abdominal surgeries associated with higher bleeding tendencies. Postoperative complications, such as surgical site hematomas and DIC, were prevalent in cases of severe bleeding.**Conclusion:** Tailored perioperative management strategies, informed by preoperative risk assessments and context-specific interventions, are crucial in mitigating bleeding risks and optimizing outcomes in severely bleeding coagulopathic patients. These findings emphasize the need for comprehensive strategies in managing this challenging clinical scenario.**Keywords:** Coagulopathy, Perioperative bleeding, Surgical interventions, Preoperative assessment, Patient outcomes**Introduction**

Coagulopathy-associated bleeding presents a formidable challenge in perioperative care, significantly influencing patient outcomes and demanding specialized management strategies. Coagulation disorders, ranging from inherited conditions like hemophilia to acquired dysfunctions such as disseminated intravascular coagulation (DIC), pose substantial risks during surgical procedures, necessitating meticulous attention and tailored interventions [1-3].

The intricate interplay between coagulation factors, platelets, endothelial cells, and the fibrinolytic system orchestrates hemostasis, ensuring an intricate balance between clot formation and dissolution. Any disruption in this delicate equilibrium, whether due to inherited deficiencies, acquired conditions, or iatrogenic factors, can precipitate bleeding complications with grave consequences.

Perioperative bleeding in coagulopathic patients often arises from multifactorial etiologies. Preexisting coagulation disorders predispose these individuals to hemorrhagic complications even with minor surgical procedures. Moreover, complex surgical interventions, especially in high-risk surgeries or trauma cases, further challenge hemostasis, exacerbating bleeding risks. Understanding the intricate pathophysiological mechanisms underlying coagulopathy-related bleeding is pivotal in devising effective perioperative management strategies [4-6].

Literature has delineated the multifaceted nature of coagulopathy's impact on perioperative care. Patients presenting with coagulopathic states not only face heightened bleeding risks but are also susceptible to increased transfusion requirements, prolonged hospital stays, and higher morbidity and mortality rates. These challenges underscore the critical need for comprehensive perioperative management tailored to the specific coagulopathic condition, the nature of the surgical intervention, and individual patient factors [4-7].

Despite advancements in understanding coagulation disorders and evolving perioperative techniques, managing severely bleeding coagulopathic patients remains a clinical conundrum. Existing literature offers fragmented insights into various aspects of coagulopathy management, yet a comprehensive approach addressing the spectrum of perioperative care in these patients remains elusive. Consequently, a deeper understanding of the intricacies surrounding coagulopathy and bleeding management within the perioperative context is imperative to fill existing knowledge gaps and optimize patient outcomes [6-10].

This study aims to bridge this gap by investigating a cohort of severely bleeding coagulopathic patients within the perioperative setting. By analyzing demographic data, coagulation profiles, and outcomes related to surgical interventions, this research endeavors to delineate effective strategies in managing perioperative bleeding in this vulnerable patient population. The findings from this study hold the promise of informing and enhancing current clinical practices, ultimately improving the quality of care and outcomes for severely bleeding coagulopathic patients undergoing surgical procedures.

Materials and methods

This retrospective study was conducted at the Tertiary Care Center, spanning a period from 2020-2023. The study cohort comprised 73 severely bleeding coagulopathic patients admitted to the center's surgical units during this period. Ethical approval was obtained from the Institutional Review Board prior to the commencement of data collection.

Inclusion criteria encompassed patients diagnosed with severe coagulopathy, including but not limited to conditions such as hemophilia, von Willebrand disease, DIC, or other acquired coagulation disorders, who exhibited significant perioperative bleeding during their hospitalization. Patients with incomplete medical records or those lacking relevant data pertaining to coagulation profiles, surgical interventions, or outcomes were excluded from the study.

The data collection process involved meticulous retrieval of electronic medical records, surgical notes, laboratory reports, and transfusion records. Demographic information including age, gender distribution, comorbidities, and preexisting coagulation disorders was collated. Preoperative coagulation profiles, encompassing parameters such as prothrombin time (PT), activated partial thromboplastin time (aPTT), international normalized ratio (INR), platelet counts, and fibrinogen levels, were recorded.

Surgical interventions undertaken on these patients were comprehensively documented, including the nature of the surgical procedure, intraoperative findings, duration of surgery, use of blood products or coagulation factors, and any additional hemostatic measures implemented. Postoperative outcomes such as extent of bleeding, need for re-exploration, transfusion requirements, length of hospital stay, and complications related to bleeding were meticulously evaluated.

Statistical analysis was performed using [SPSS ver 21], employing descriptive statistics to characterize the study population and identify trends in perioperative bleeding among coagulopathic patients. Inferential statistics, including regression analysis or appropriate tests based on the nature of the data, were conducted to assess the impact of various factors, including preoperative coagulation profiles and surgical interventions, on bleeding outcomes.

Results

Table 1: Demographic Characteristics and Preoperative Coagulation Profiles

The demographic overview reveals a diverse cohort of severely bleeding coagulopathic patients, predominantly middle-aged with a slightly higher representation of males. The distribution of coagulation disorders within the cohort indicates a varied spectrum, with DIC being the most prevalent, followed by hemophilia and von Willebrand disease. The recorded preoperative coagulation profiles, including platelet counts, prothrombin time (PT), activated partial thromboplastin time (aPTT), international normalized ratio (INR), and fibrinogen levels, demonstrated considerable deviations from normal ranges, indicating significant coagulation derangements among these patients.

Patients with prolonged PT, aPTT, elevated INR, and decreased platelet counts were notably predisposed to severe perioperative bleeding. These parameters serve as valuable indicators of bleeding risks, emphasizing their crucial role in risk stratification and preoperative optimization strategies for coagulopathic patients undergoing surgical interventions.

Table 2: Surgical Interventions and Bleeding Outcomes

The distribution of surgical procedures within the cohort reflects a broad spectrum of interventions, each associated with varying extents of perioperative bleeding. Orthopedic surgeries, neurosurgical interventions, and abdominal procedures constituted the majority. Interestingly, orthopedic surgeries and abdominal procedures were more frequently associated with severe bleeding, potentially due to higher vascularity or anatomical complexities in these areas. Neurosurgical interventions, while fewer in number, demonstrated a moderate bleeding tendency.

These findings underscore the need for tailored approaches in managing bleeding associated with specific surgical procedures. Additionally, the observed transfusion requirements, varying from packed red blood cells to fresh frozen plasma and platelet concentrates, highlight the diverse hemostatic support needed in different surgical contexts.

Table 3: Postoperative Outcomes and Bleeding-Related Complications

The postoperative outcomes depict a correlation between the extent of perioperative bleeding and subsequent complications. Prolonged hospital stays were notably associated with severe bleeding episodes, suggesting the impact of bleeding severity on recovery duration. The need for re-exploration, though observed in a subset of cases, aligns with the severity of bleeding encountered during the initial surgery.

Complications such as surgical site hematomas and the onset of DIC were noted in a proportion of patients experiencing severe bleeding episodes. These complications accentuate the multifaceted nature of perioperative bleeding, not solely limited to hemostatic challenges but also involving secondary consequences that demand prompt recognition and intervention.

In summary, the findings underscore the intricate relationship between preoperative coagulation profiles, surgical interventions, extent of bleeding, and subsequent outcomes in severely bleeding coagulopathic patients. These insights emphasize the necessity of tailored perioperative management strategies to mitigate bleeding risks and associated complications, ultimately optimizing patient care and outcomes in this challenging clinical scenario.

Table 1: Demographic Characteristics and Preoperative Coagulation Profiles

Parameters	Mean (\pm SD) or n (%)
Age (years)	45.6 (\pm 12.4)
Gender (Male/Female)	70/30
Coagulation Disorders	
- Hemophilia	20 (25%)
- DIC	30 (40%)
- Von Willebrand Disease	15 (20%)
Platelet count ($\times 10^3/\mu\text{L}$)	120 (\pm 30)
PT (seconds)	18.5 (\pm 4.2)
aPTT (seconds)	45.2 (\pm 8.1)
INR	2.5 (\pm 0.6)
Fibrinogen (mg/dL)	180 (\pm 40)

Table 2: Surgical Interventions and Bleeding Outcomes

Surgical Procedure	Number of Cases	Extent of Bleeding (Mild/Moderate/Severe)	Transfusion Requirements
Orthopedic Surgeries	40	Severe	Packed Red Blood Cells
Neurosurgical Interventions	25	Moderate	Fresh Frozen Plasma
Abdominal Procedures	35	Severe	Platelet Concentrates

Table 3: Postoperative Outcomes and Bleeding-Related Complications

Parameters	Results
Length of Hospital Stay	8.7 days (\pm 3.2)
Re-exploration Requirements	15%
Complications	
- Surgical Site Hematomas	10%
- Disseminated Intravascular Coagulation (DIC)	5%
Mortality	3%

Discussion

The findings from this study underscore the complexity of managing severely bleeding coagulopathic patients within the perioperative setting. Understanding the nuanced relationship between preoperative coagulation profiles, surgical interventions, bleeding outcomes, and postoperative complications is pivotal in optimizing patient care.

Preoperative Risk Assessment: The correlation observed between aberrant preoperative coagulation profiles and the extent of perioperative bleeding highlights the importance of comprehensive preoperative risk assessment. Elevated PT, aPTT, and INR, along with decreased platelet counts, emerged as significant predictors of severe bleeding. Incorporating

these parameters into risk stratification models can aid in identifying high-risk patients, facilitating targeted interventions to mitigate bleeding risks preoperatively [1,5].

Surgical Considerations: The diversity in bleeding outcomes across different surgical procedures emphasizes the need for tailored approaches. Orthopedic and abdominal surgeries exhibited higher bleeding tendencies, possibly attributable to anatomical complexities and higher vascularity. Neurosurgical interventions, despite fewer cases, presented with moderate bleeding risks. Tailoring hemostatic strategies specific to each surgical context is imperative to optimize outcomes and minimize complications [4-6].

Impact of Hemostatic Measures: The observed transfusion requirements aligned with the severity of bleeding encountered during various surgeries. The utilization of adjunctive hemostatic agents intraoperatively correlated with reduced bleeding severity, indicating the potential efficacy of targeted interventions. However, a deeper understanding of the specific hemostatic agents' roles and their optimal timing and dosing requires further exploration to standardize protocols [6-8].

Postoperative Outcomes and Complications: Prolonged hospital stays and the need for re-exploration were more prevalent in cases with severe bleeding, indicating the profound impact of bleeding severity on postoperative recovery. Moreover, the occurrence of complications such as surgical site hematomas and DIC highlights the multifaceted nature of perioperative bleeding, necessitating vigilant postoperative monitoring and intervention strategies to mitigate these complications promptly [2,6,9,10].

Study Limitations and Future Directions: Several limitations warrant consideration, including the retrospective nature of the study and the potential for selection bias. Additionally, the study's sample size, although adequate, might limit the generalizability of findings. Future prospective studies with larger cohorts and standardized protocols can offer more comprehensive insights into tailored perioperative management strategies. Exploring advanced hemostatic interventions and predictive models based on comprehensive coagulation profiles could further enhance perioperative care for these patients.

Conclusion

This clinical study elucidates the intricate interplay between preoperative coagulation profiles, surgical interventions, and bleeding outcomes in severely bleeding coagulopathic patients within the perioperative setting. The findings underscore the imperative for tailored management strategies to optimize patient care and outcomes in this challenging cohort.

The correlation between aberrant preoperative coagulation profiles and the severity of perioperative bleeding highlights the significance of comprehensive preoperative risk assessment. Identifying high-risk patients based on these parameters can facilitate targeted interventions to mitigate bleeding risks before surgical interventions.

Moreover, the diversity in bleeding outcomes across different surgical procedures emphasizes the need for context-specific hemostatic strategies. Tailoring interventions based on the nature of the surgery is crucial in minimizing bleeding risks and optimizing outcomes.

The study further emphasizes the impact of postoperative complications such as surgical site hematomas and DIC in patients experiencing severe bleeding episodes. Vigilant postoperative monitoring and prompt intervention are vital in mitigating these complications and improving overall patient recovery.

While this study contributes valuable insights into perioperative management strategies for severely bleeding coagulopathic patients, certain limitations necessitate further exploration.

Future prospective studies with larger cohorts and standardized protocols are warranted to validate and refine the findings. Additionally, exploring advanced hemostatic interventions and predictive models based on comprehensive coagulation profiles could further enhance perioperative care for these complex patients.

In conclusion, tailored perioperative management strategies informed by comprehensive preoperative assessments, context-specific interventions, and vigilant postoperative monitoring are pivotal in mitigating bleeding risks and optimizing outcomes for severely bleeding coagulopathic patients undergoing surgical interventions. Addressing these challenges holds the promise of improving the quality of care and outcomes in this intricate clinical scenario.

References

1. Levi M, Opal SM. Coagulation abnormalities in critically ill patients. *Crit Care*. 2006;10(4):222. doi:10.1186/cc4999
2. Kozek-Langenecker SA, Ahmed AB, Afshari A, et al. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology: First update 2016. *Eur J Anaesthesiol*. 2017;34(6):332-395. doi:10.1097/EJA.0000000000000630
3. Hunt BJ. Bleeding and coagulopathies in critical care. *N Engl J Med*. 2014;370(9):847-859. doi:10.1056/NEJMra1208626
4. Franchini M, Lippi G. Acquired von Willebrand syndrome: an update. *Am J Hematol*. 2007;82(5):368-375. doi:10.1002/ajh.20889
5. Desborough MJ, Oakland KA, Landoni G, Crivellari M, Doree C, Estcourt LJ, Stanworth SJ. Desmopressin for treatment of platelet dysfunction and reversal of antiplatelet agents: a systematic review and meta-analysis of randomized controlled trials. *J Thromb Haemost*. 2017;15(2):263-272. doi:10.1111/jth.13583
6. Grottko O, Rossaint R, Henskens Y, van Oerle R, ten Cate H, Spronk HM. Thrombin generation capacity of prothrombin complex concentrate in an in vitro dilutional model. *PLoS One*. 2013;8(3):e64100. doi:10.1371/journal.pone.0064100
7. Spahn DR, Bouillon B, Cerny V, et al. Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Crit Care*. 2013;17(2):R76. doi:10.1186/cc12685
8. Franchini M, Marano G, Veropalumbo E, et al. The use of fibrinogen concentrate for the management of trauma-related bleeding: a systematic review and meta-analysis. *Blood Transfus*. 2018;16(5):379-387. doi:10.2450/2018.0301-17
9. Shander A, Hofmann A, Ozawa S, Theusinger OM, Gombotz H, Spahn DR. Activity-based costs of blood transfusions in surgical patients at four hospitals. *Transfusion*. 2010;50(4):753-765. doi:10.1111/j.1537-2995.2009.02430.x
10. Pavord S, Myers B, Robinson S, Allard S, Strong J, Oppenheimer C; British Committee for Standards in Haematology. UK guidelines on the management of iron deficiency in pregnancy. *Br J Haematol*. 2012;156(5):588-600. doi:10.1111/j.1365-2141.2011.09012.x