

The effect of IASTM technique on the pain and range of motion in patients with low back pain in patients of Rama Medical College Hospital.

Dr. Gulwish Sadique¹ Dr. Manjit Kumar¹ Dr. Minhaj Tahir¹

¹ Assistant Professor, Rama Institute of Paramedical Sciences, Rama University, Kanpur

Abstract

[Purpose] Clinicians have reported the effects of various instrument assisted soft tissue mobilization (IASTM) in patients of Rama Medical College Hospital. The purpose of this study was to investigate the effects of the IASTM technique and general exercise on pain and range of motion (ROM) in patients with LBP.

[Subjects and Methods] 30 individuals with LBP participated in the study (IASTM technique: 15; Control: 15). Before and after the 4-week intervention program, pain was assessed using a visual analog scale (VAS). Lumbar ROM was measured using a smartphone. The main effects and interaction were analyzed by two-way repeated ANOVA.

[Results] A significant time-by-group interaction was observed for the VAS and ROM. A post hoc paired t-test showed that pain decreased significantly post intervention within the IASTM group. The lumbar ROM significantly increased post-intervention in both groups.

[Conclusion] The IASTM technique and general exercise resulted in pain relief and increased ROM. However, the IASTM group showed significantly increased VAS and ROM more than control group. These findings suggest that the IASTM technique can be useful as a pain decrease and ROM increase for patients with LBP.

Key words: low back pain, IASTM technique, Range of motion.

INTRODUCTION

LBP patients when bending forward to going back because of the balance of the posterior structures such as ligaments and erector spinae¹. LBP patients had neuromuscular problem or a problem of internal muscles, muscle weakness, pain due to shortening occurs complain². A significant difference in muscular activities of erector spinae between the groups were obtained when returning to the erect position from the maximum flexion. Moreover, time lag between trunk and hip movement was much greater in patients of Rama medical college and hospital than in healthy subjects. This study demonstrated that neuromuscular coordination between trunk and hip could be abnormal in patients with LBP³. Instrument assisted soft tissue mobilization is a new range of tool which enables clinicians to efficiently locate and treat individuals diagnosed with soft tissue dysfunction. IASTM is a procedure that is rapidly growing in popularity due to its effectiveness and efficiency while remaining non-invasive, with its own indications and limitations. Gehlsen et al.⁴ investigated the effects of 3 separate IASTM pressures on rat Achilles tendons. They concluded that fibroblast production is directly proportional to the magnitude of IASTM pressure used by the clinician. Davidson et al.⁵ supported Gehlsen et al.⁴ by concluding that intramuscular (IM) significantly increased fibroblast production in rat Achilles tendons by using electron microscopy to analyze tissue samples following IM application. Davidson et al.⁵ found morphologic changes in the rough endoplasmic reticulum following IM application. Thus, indicating micro trauma to damaged tissues, resulting in an acute fibroblast response⁵. The results of this study indicate that an application of IASTM to the posterior shoulder provides acute improvements in both glenohumeral horizontal adduction range of motion (ROM) and internal rotation ROM among baseball players⁶. In the present study, we determined the changes in the pain and ROM of back muscles after therapy using the IASTM technique in patients with LBP.

We hypothesized that IASTM technique would increase ROM and decrease pain more than the general exercise.

SUBJECTS AND METHODS

This study involved 30 patients of Rama Medical College Hospital (17 females, 13 males) with low back pain. Inclusion criteria included onset of low back pain of less than 12 weeks, LBP (>90 days at time of enrolment⁷). Exclusion criteria were a history of back surgery⁸, patients who had spinal fracture within six months and spinal tumor or other malignancy and medicine for psychiatric disorder and who had possibility of exaggerated complaints due to automobile or accident insurance⁷. All of the participants read and signed an informed consent. Control group (n=15) age 33.0 ± 9.9 , height 165.6 ± 7.9 , body weight 61.8 ± 12.9 , and VAS score 48.9 ± 14.6 . Graston group (n=15) age 40.6 ± 14.6 , height 169.3 ± 10.2 , body weight 63.9 ± 15.6 , and VAS score 50.6 ± 12.7 . To measure lumbar ROM, hip ROM inclinometer application, Android 4.0 phone (Samsung Galaxy S3, SHV-E210S) was used. Data were collected using an Android 4.0 phone ROM was recorded. The intra class correlation coefficients ranged from 0.82 to 0.902).

The IASTM technique was used for the group for intervention. Pain level was evaluated using visual analog scale (VAS; 100 mm)^{1, 8, 9}. Zero indicated no pain and 100 mm indicate the worst pain level. The VAS score of 17 mm is reported as minimal clinically important difference¹⁰). Lumbar, hip ROM was measured with an inclinometer application. To measure lumbar flexion/extension, the neutral position, with feet 15 cm apart and heels aligned with floor tape. Smartphone placement were used at the T12–L1 between interspinous junction and S1 Tubercle¹¹). The patient was asked to maximally flex/extend, keeping the knees straight, at which points readings were taken from inclinometer application. The method purpose is to record true lumbar flexion/extension without sacral involvement¹).

To measure lumbar lateral bending, smartphone placed at T12–L1 between over the sacrum according. The patient in full lateral bending right/left. Hip flexion ROM was measured with participants in the supine position smartphone was firmly attached with adhesive tape to the anterior aspect of the thigh, two centimeters proximal to the superior pole of the patella. The reading on the smartphone held against the skin laterally at the midpoint of the thigh (determined as midway between the lateral femoral condyle and greater trochanter) with the leg relaxed in neutral hip rotation, were recorded¹³). Maximal active hip flexion was then performed with the knee in flexion and recorded at the point at which a pain was experienced. This position was then maintained for three seconds to allow the smartphone to record the measurement¹³). We measured each motion in three trials and recorded the average. The IASTM group performed IASTM using during 4 weeks: posterior fascia, sacrum, hip lateral rotators, and hamstring bilaterally and general exercise. In the first, subjects were asked to kneel directly on the bed. The posterior fascia IASTM was microtrauma to stimulation lumbar posterior muscle erector spinae (iliocostalis, longissimus), and multifidus. The sacrum IASTM, subjects were asked to kneel directly on the bed. The hip lateral rotators IASTM was applied to kneeling prone position with knee and hip flexion sidelying position at gluteus maximus and gluteus medius. The hamstring bilaterally IASTM was applied to prone position at biceps femoris, semitendinosus, and semimembranosus. The IASTM treatment was applied for approximately 20-seconds in a direction parallel to the muscle fibers being treated with the instrument at a 45° angle. Followed immediately by treating the muscles in a direction perpendicular to the muscle fibers with the instrument at a 45° angle for an additional 20-seconds, resulting in a total treatment time of approximately 40 seconds. The control group was applied general exercise. General exercise was applied with stretching exercises and stationary bicycling

for 10–15 minutes^{14, 15}). Three sets of fifteen repetitions were performed, with rest times of 1 minute between sets during 4 weeks. We expressed the mean and standard deviation (SD) of the VAS scores and ROM. Statistical analyses were performed with the SPSS software (ver. 22.0 for Windows; SPSS Inc., Chicago, IL, USA). A two-way repeated-measures analysis of variance (ANOVA) was used to determine the main effect and any interaction between the VAS score and ROM. The within-group factor was time (pre-intervention vs. post-intervention) and the between-group factor was group (IASTM vs. Control). A p-value<0.05 was considered statistically significant. If a significant main effect or time-by-group interaction effect was detected, a post hoc independent and paired t-test was used.

RESULTS

Significant main effects of time were observed in pain. VAS scores significantly decreased post-intervention versus pre intervention in IASTM groups (Table 1; IASTM: 25.5 ± 7.3 mm vs. 50.6 ± 12.8 mm, p<0.001; Control: 44.6 ± 12.9 vs. 48.9 ± 14.6, p=0.334). Significant main effects of time were observed in ROM. ROM of lumbar flexion significantly increased post-intervention

Table 1. Change in pain between pre- and post-intervention post (N=30)

Variable	Group	Pre-	Post-
		intervention	intervention
		Mean ± SD	Mean ± SD
VAS (mm)	Graston (n=15)	50.6 ± 12.7	25.5 ± 7.3*
	Control (n=15)	48.9 ± 14.6	44.6 ± 12.9

*p<0.05. VAS: visual analog scale.

Table 2. Change in range of motion between pre- and intervention (N=30)

Variable	Group	Pre-	Post-
		intervention	intervention
		Mean ± SD	Mean ± SD
Lumbar flexion	IASTM (n=15)	74.5 ± 14.1	89.3 ± 10.5*
	Control (n=15)	78.2 ± 8.7	75.5 ± 17.5*
Lumbar extension	IASTM (n=15)	13.0 ± 3.2	19.8 ± 2.4*
	Control (n=15)	13.3 ± 2.4	14.6 ± 2.3*
Lumbar lateral bending (Rt)	IASTM (n=15)	22.1 ± 4.4	25.6 ± 4.7*
	Control (n=15)	22.9 ± 3.3	23.6 ± 3.3*
Lumbar lateral bending (Lt)	IASTM (n=15)	21.3 ± 4.2	25.7 ± 4.6*
	Control (n=15)	23.2 ± 3.3	24.2 ± 3.3*
Hip flexion	IASTM (n=15)	110.2 ± 6.6	118.1 ± 8.1*
	Control (n=15)	110.6 ± 6.8	111.8 ± 5.5*

*p<0.05

versus pre-intervention in both groups (Table 2; IASTM: 89.3 ± 10.5° vs. 74.5 ± 14.2°, p<0.001; Control: 75.5 ± 17.5° vs. 78.3 ± 8.8°, p=0.492). ROM of lumbar extension significantly increased post-intervention versus pre-intervention in both groups (Table 2; IASTM: 19.8 ± 2.42° vs. 13.0 ± 3.2°, p<0.001; Control: 14.6 ± 2.3° vs. 13.3 ± 2.4°, p=0.026). ROM of lateral bending (right) significantly increased post-intervention versus pre-intervention in both groups (Table 2; IASTM: 25.6 ± 4.7° vs. 23.6 ± 3.3°, p<0.001; Control: 23.6 ± 3.3° vs. 22.9 ± 3.3°, p=0.002). ROM of lateral bending (left) significantly increased post-intervention versus post-intervention in both groups (IASTM: 25.7 ± 4.6° vs. 21.3 ± 4.2°, p<0.001; Control: 24.2 ± 3.3° vs. 23.2 ± 3.3°, p=0.014). ROM of hip flexion significantly increased post-intervention versus post-intervention in both groups (IASTM: 118.1 ± 8.1° vs. 110.2 ± 6.6°, p<0.001; Control: 111.8 ± 5.5° vs. 110.6 ± 6.8°, p=0.021). There were significant differences in lumbar and hip ROM between groups.

DISCUSSION

Many researchers have investigated the effects of soft tissue mobilization in patients^{9, 15-17}). However, few studies have investigated the effect of soft tissue mobilization on ROM, pain in patients with LBP. We investigated the effects of the IASTM technique compare with general exercise in patients with LBP. Our results show that the IASTM technique decreased pain and increased ROM in patients with LBP. These results confirm that both methods lead to pain relief and increased ROM. These findings indicate that the IASTM technique could be recommended in the rehabilitation of patients with LBP. The IASTM group showed improvements decreased pain and increased ROM. In this study, pain significantly decreased in IASTM group. Our results are consistent with other reports^{15, 18}). Previous studies explained that pain decreases could be affected erector spinae activity^{1, 19, 20} and back muscle activity was decrease^{21, 22}.

Our results demonstrated that after performing the IASTM technique pain significantly decreased. Thus, IASTM technique may help to decrease lumbar pain in patients with LBP. IASTM technique more than general exercise under these conditions may decrease muscle activities, leading to decreased back pain. A VAS score of 17 mm is the minimal clinically important difference¹⁰). Mean differences in VAS score reduced pre- and post-intervention were 25.1 mm in the IASTM group and 4.3 mm in the control group. We propose that both the IASTM technique compare with general exercise contributed to decrease pain in patients with LBP. Active lumbar flexion, extension, lateral bending and hip flexion were increased both in the IASTM technique and control group. IASTM technique and general exercises are used to increase ROM and may do nothing more; a meta-analysis indicated that pain modality did not significantly increase ROM²².

This was because the IASTM technique helped to prevent muscle atrophy and restored muscle balance and general exercise included stretching for improving ROM. Thus, the hypothesis of the present study was confirmed, as IASTM technique increased lumbar and hip ROM more significantly than the general exercise. However, both the IASTM and control groups showed significantly increased lumbar and hip ROM after the interventions. Consequently, the IASTM technique may be a useful intervention for ROM of lumbar and hip. The present study had several limitations. First, it only investigated changes in pain, ROM after the IASTM technique, and did not measure the muscles activity. Thus, further studies are required to investigate changes in back muscle activity after the IASTM technique. Second, the duration of intervention was short. Usually, a 6–8 week or longer intervention period has been used because the most significant progress may be made after at least 6 weeks^{23, 24}). However, what is the optimal exercise period remains unclear²⁵. Our results showed that after 4 weeks, the IASTM technique significantly increased ROM and decreased pain. Further research is needed to investigate the long-term effects of the IASTM technique on pain muscle activity and ROM in patients with LBP.

Table 1. Change in pain between pre- and post-intervention

(N=30)

Variable Group

Pre intervention

Post intervention

Mean ± SD Mean ± SD

VAS

(mm)

IASTM (n=15) 50.6 ± 12.7 25.5 ± 7.3*

Control (n=15) 48.9 ± 14.6 44.6 ± 12.9

*p<0.05. VAS: visual analog scale.

Table 2. Change in range of motion between pre- and post-intervention

(N=30)

Variable Group

Pre intervention

Post intervention

Mean \pm SD Mean \pm SD

Lumbar

flexion

IASTM (n=15) 74.5 \pm 14.1 89.3 \pm 10.5*

Control (n=15) 78.2 \pm 8.7 75.5 \pm 17.5*

Lumbar

extension

IASTM (n=15) 13.0 \pm 3.2 19.8 \pm 2.4*

Control (n=15) 13.3 \pm 2.4 14.6 \pm 2.3*

Lumbar lateral

bending (Rt)

IASTM (n=15) 22.1 \pm 4.4 25.6 \pm 4.7*

Control (n=15) 22.9 \pm 3.3 23.6 \pm 3.3*

Lumbar lateral

bending (Lt)

IASTM (n=15) 21.3 \pm 4.2 25.7 \pm 4.6*

Control (n=15) 23.2 \pm 3.3 24.2 \pm 3.3*

Hip flexion

IASTM (n=15) 110.2 \pm 6.6 118.1 \pm 8.1*

Control (n=15) 110.6 \pm 6.8 111.8 \pm 5.5*

*p<0.05

We investigated the effects of the IASTM technique on pain and ROM in patients with low back pain. Both the IASTM technique and general exercise induced increased ROM. However, only the IASTM group showed more pain relief and increased ROM. These findings suggest that the IASTM technique can be useful as a pain relief and ROM increasing program for patients with LBP.

REFERENCES

- 1) Dubois JD, Piché M, Cantin V, et al.: Effect of experimental low back pain on neuromuscular control of the trunk in healthy volunteers and patients with chronic low back pain. *J Electromyogr Kinesiol*, 2011, 21: 774–781. [Medline] [CrossRef]
- 2) Hartvigsen L, Kongsted A, Hestbaek L: Clinical examination findings as prognostic factors in low back pain: a systematic review of the literature. *Chiropr Man Therap*, 2015, 23: 13. [Medline] [CrossRef]
- 3) Scott IR, Vaughan AR, Hall J: Swiss ball enhances lumbar multifidus activity in chronic low back pain. *Phys Ther Sport*, 2015, 16: 40–44. [Medline] [Cross-Ref]
- 4) Gehlsen GM, Ganion LR, Helfst R: Fibroblast responses to variation in soft tissue mobilization pressure. *Med Sci Sports Exerc*, 1999, 31: 531–535. [Medline] [CrossRef]
- 5) Davidson CJ, Ganion LR, Gehlsen GM, et al.: Rat tendon morphologic and functional changes resulting from soft tissue mobilization. *Med Sci Sports Exerc*, 1997, 29: 313–319. [Medline] [CrossRef]
- 6) Laudner K, Compton BD, McLoda TA, et al.: Acute effects of instrument assisted soft tissue mobilization for improving posterior shoulder range of motion in collegiate baseball players. *Int J Sports Phys Ther*, 2014, 9: 1–7. [Medline]
- 7) Lee SW, Kim SY: Effects of hip exercises for chronic low-back pain patients with lumbar instability. *J Phys Ther Sci*, 2015, 27: 345–348. [Medline] [CrossRef]
- 8) Deckers K, De Smedt K, van Buyten JP, et al.: Chronic low back pain: restoration of dynamic stability. *Neuromodulation*, 2015, 18: 478–486, discussion 486. [Medline] [CrossRef]
- 9) Han GS, Cho MH, Nam KT, et al.: The effects on muscle strength and visual analog scale pain of aquatic therapy for individuals with low back pain. *J Phys Ther Sci*, 2011, 23: 57–60. [CrossRef]
- 10) Mark HSM, AU TTS, Choi YF, et al.: The minimum clinically significant difference in visual analogue scale pain score in a local emergency setting. *Hong Kong J Emerg Med*, 2009, 16: 233–236.
- 11) Breum J, Wiberg J, Bolton JE: Reliability and concurrent validity of the BROM II for measuring lumbar mobility. *J Manipulative Physiol Ther*, 1995, 18: 497–502. [Medline]
- 12) Keeley J, Mayer TG, Cox R, et al.: Quantification of lumbar function. Part 5: Reliability of range-of-motion measures in the sagittal plane and an in vivo torso rotation measurement technique. *Spine*, 1986, 11: 31–35. [Medline] [CrossRef]
- 13) Charlton PC, Mentiplay BF, Pua YH, et al.: Reliability and concurrent validity of a Smartphone, bubble inclinometer and motion analysis system for measurement of hip joint range of motion. *J Sci Med Sport*, 2015, 18: 262–267. [Medline] [CrossRef]
- 14) Ludewig PM, Hoff MS, Osowski EE, et al.: Relative balance of serratus anterior and upper trapezius muscle activity during push-up exercises. *Am J Sports Med*, 2004, 32: 484–493. [Medline] [CrossRef]
- 15) Chatchawan U, Jupamatangb U, Chanchitc S, et al.: Immediate effects of dynamic sitting exercise on the lower back mobility of sedentary young adults. *J Phys Ther Sci*, 2015, 27: 3359–3363. [Medline] [CrossRef]
- 16) Burke J, Buchberger DJ, Carey-Loghmani MT, et al.: A pilot study comparing two manual therapy interventions for carpal tunnel syndrome. *J Manipulative*

Physiol Ther, 2007, 30: 50–61. [Medline] [CrossRef]

17) Christenson RE: Effectiveness of specific soft tissue mobilizations for the management of Achilles tendinosis: single case study—experimental design. Man

Ther, 2007, 12: 63–71. [Medline] [CrossRef]

18) Hunter G: Specific soft tissue mobilization in the management of soft tissue dysfunction. Man Ther, 1998, 3: 2–11. [Medline] [CrossRef]

19) Aspegren D, Hyde T, Miller M: Conservative treatment of a female collegiate volleyball player with costochondritis. J Manipulative Physiol Ther, 2007, 30:

321–325. [Medline] [CrossRef]

20) Lee CW, Hwangbo K, Lee IS: The effects of combination patterns of proprioceptive neuromuscular facilitation and ball exercise on pain and muscle activity

of chronic low back pain patients. J Phys Ther Sci, 2014, 26: 93–96. [Medline] [CrossRef]

21) Saner J, Kool J, Sieben JM, et al.: A tailored exercise program versus general exercise for a subgroup of patients with low back pain and movement control

impairment: a randomised controlled trial with one-year follow-up. Man Ther, 2015, 20: 672–679. [Medline] [CrossRef]

22) Cho HY, Kim EH, Kim J: Effects of the CORE exercise program on pain and active range of motion in patients with chronic low back pain. J Phys Ther Sci,

2014, 26: 1237–1240. [Medline] [CrossRef]

23) Hammer WI, Pfefer MT: Treatment of a case of subacute lumbar compartment syndrome using the Graston technique. J Manipulative Physiol Ther, 2005, 28:

199–204. [Medline] [CrossRef]

24) Blanchette MA, Normand MC: Augmented soft tissue mobilization vs natural history in the treatment of lateral epicondylitis: a pilot study. J Manipulative

Physiol Ther, 2011, 34: 123–130. [Medline] [CrossRef]

25) Arokoski JP, Valta T, Kankaanpää M, et al.: Activation of lumbar paraspinal and abdominal muscles during therapeutic exercises in chronic low back pain

patients. Arch Phys Med Rehabil, 2004, 85: 823–832. [Medline] [CrossRef]