

**Title of the article- THE INCIDENCE OF RETINOPATHY OF
PREMATURITY AT A TERTIARY HOSPITAL**

Type of article- Original

Authors Name and affiliations-

1. Dr Nikita Bilania

Postgraduate student JR 2

Dept of Ophthalmology, VCSGGIMSR, Srinagar garhwal

2. Dr Jigyasa Singh

Postgraduate student JR 3

Dept of Ophthalmology, VCSGGIMSR, Srinagar garhwal

3. Dr Dharmendra Kumar

Senior resident

Dept of ophthalmology VCSGGIMSR srinagar Garhwal

4. Dr Achyut N Pandey

Associate professor

Dept of ophthalmology VCSGGIMSR srinagar Garhwal

Corresponding author

Dr Achyut N Pandey

Associate Professor

Dept of Ophthalmology, VCSGGIMSR, Srinagar garhwal

9456792024

achyutpandey@gmail.com

Financial Interest

NIL

ABSTRACT:

Purpose: To study the incidence and risk factors of retinopathy of prematurity (ROP).

Methods: A prospective observational study of premature babies admitted in NICU from December 2021 to December 2022 with a gestational age of 32 weeks or less at birth and a birth weight of 1500 g or less was made. Sick infants were included even if they were older and heavier. ROP screening was done in 200 cases from the fourth postnatal week and was followed up.

Results: The incidence of ROP was 19.5%. Most common maternal risk factor was pregnancy induced hypertension (17.9%). Low birth weight (LBW) and respiratory distress syndrome (RDS) (89.7%) were the most common risk factors in infants.

Conclusion: In our study the most common risk factors were LBW and RDS.

INTRODUCTION:

Retinopathy of prematurity (ROP) is characterised by abnormal development of retinal vasculature. It is an important and preventable cause of childhood blindness¹. 50,000 cases of childhood blindness in the world every year are estimated to be caused by ROP². Due to advancement in the health care system, survival rates of extremely premature babies have increased. This has led to an increase in the incidence of ROP in developing countries³. Previously, a major risk factor for development of ROP was thought to be use of high concentration of supplemental oxygen therapy^{4,5}. However, ROP has been observed in patients without oxygen therapy⁶. Also, low birth weight (BW) and low gestational age (GA) are well-known risk factors for ROP^{7,8}.

The criteria for ROP screening has been well determined in industrialised countries but since the blindness from ROP varies from region to region, a single screening program cannot suit all regions as seen in various studies⁹.

PURPOSE:

To study the incidence and risk factors of retinopathy of prematurity (ROP) at a tertiary hospital.

METHODS:

This is a prospective observational study which included premature babies who were referred for ROP screening, admitted in NICU from December 2021 to December 2022, with a gestational age of 32 weeks or less at birth and a birth weight of 1500 g or less. Babies >32 weeks and >1500 grams were screened if the babies were sick as per

paediatricians advice. ROP screening was done from the fourth postnatal week and within four weeks if the baby was ≤ 30 weeks and was followed up accordingly.

Data on birth weight, gestational age, post conceptional age was noted. Maternal risk factors like (pregnancy induced hypertension, gestational diabetes) and baby risk factors like (respiratory distress syndrome, sepsis) were also noted in the history.

Babies in whom media opacity (not related to ROP) precluding fundus visualization disease, loss to follow-up were excluded from the study.

The pupils were dilated by application of tropicamide 0.5% and phenylephrine 1%, and the patients were examined by a single retina specialist using a binocular indirect ophthalmoscope with 20 D lens. A sterile lid speculum was used.

ROP was classified according to the international classification of ROP and follow-up schedules were designed in accordance with suggestions of the American Academy of Pediatrics, American Academy of Ophthalmology and American Association for Pediatric Ophthalmology and Strabismus¹⁰.

RESULTS:

During the period of 1 year, 200 babies were screened. Male babies screened were 110, and female babies were 90. Gestational age (GA) of babies ranged from 26 weeks-38 weeks (meanGA-33wks). Birth weight (BW) of the babies range from 1kg-3.5kg (meanBW-2.25kg). Incidence of ROP was 19.5% (39 babies) in our study. Different stages of ROP seen are as shown in the table1. In the babies diagnosed to have ROP the gestational age ranged from 26weeks- 37weeks (with 30% being more than 32 weeks). Birth weight ranged from 1kg-2kg (with 25.6% being more than 1.5kg). Most common maternal risk factor for ROP was pregnancy induced hypertension in 7 patients (17.9%) followed by gestational diabetes in 2 cases (5.1%). The most common neonatal risk factors for ROP was respiratory distress syndrome in 35 babies (89.7%) followed by sepsis in 5 cases (12.8%).

TABLE1:

ROP(stage)	NO. (%)
Stage 1	5 (12.8%)
Stage 2	16 (41%)
Stage 3	18 (46.2%)
Stage 4	0
Stage 5	0

DISCUSSION:

ROP is an important cause of childhood blindness, which if diagnosed and managed early can prevent permanent visual loss. The incidence of ROP in premature infants in our study was 19.5%. In developed countries, the reported incidence of ROP ranges from 19.3% to 25%^(12,13). Likewise in India, the incidence of ROP depends on the region of study and ranges from 20% to 47.3%^(14,15). In the current study the incidence of ROP is almost same as other studies performed in India. As reported by Palmer, *et al.*⁽¹⁶⁾, incidence and severity of ROP was closely related to lower birth weight and lower postconceptional age, as was seen in our study.

There are varying screening criteria described by different authors. Vinekar, *et al.*⁽¹⁷⁾ suggested that the scenario in developing countries is quite different. Larger and gestationally 'older' infants are more likely to develop ROP compared to their counterparts in Western countries. Hence, the application of Western screening guidelines for developing countries has been questioned by Jalali, *et al.*⁽¹⁸⁾. As a higher cut off limit, they recommended screening babies born at <37 weeks gestation and/or birthweight <2000g in the presence of a high sickness score, in order to prevent missing any infant with threshold ROP.

So, we feel that all babies with birthweight less than 1500g and gestation ≤ 32 weeks should be routinely screened. Infants with birthweight between 1500-2000g and gestational age more than 32 weeks should be screened at the discretion of the neonatologist, depending on other risk factors during the course of stay in the NICU.

Many risk factors have been reported to predispose to the development of ROP. In our study the most common maternal risk factor noted was pregnancy induced hypertension seen in 17.9% followed by gestational diabetes in 5.1%. Most common neonatal risk factors were respiratory distress syndrome in 89.7% followed by sepsis in 12.8%.

Oxygen therapy was not found to influence the development of ROP in our study.

CONCLUSION:

ROP screening guidelines should vary according to the region. The most common risk factors were low birth weight, lower gestational age and respiratory distress syndrome and pregnancy induced hypertension.

REFERENCES:

1. Yang CS, Chen SJ, Lee FL, Hsu WM, Liu JH. Retinopathy of prematurity: Screening, incidence and risk factors analysis. *Zhonghua Yi Xue Za Zhi (Taipei)* 2001;64:706–712.
2. Saugstad OD. Oxygen and retinopathy of prematurity. *J Perinatol.* 2006;26(Suppl1):S46–50.
3. Gilbert C. Retinopathy of prematurity: A global perspective of the epidemics, population of babies at risk and implications for control. *Early Hum Dev.* 2008;84:77–82.
4. Saugstad OD. Is oxygen more toxic than currently believed? 2001;108:1203–1205.
5. Penn JS, Henry MM, Wall PT, Tolman BL. The range of PaO₂ variation determines the severity of oxygen-induced retinopathy in newborn rats. *Invest Ophthalmol Vis Sci.* 1995;36:2063–2070.
6. Lucey JF, Dangman B. A reexamination of the role of oxygen in retrolental fibroplasia. 1984;73:82–96.
7. Mousavi SZ, Karkhaneh R, Riazi-Esfahani M, Mansouri MR, Roohipoor R, Ghalichi L, et al. Retinopathy of prematurity in infants with late retinal examination. *J Ophthalmic Vis Res.* 2009;4:24–28.
8. Fortes Filho JB, Eckert GU, Procianoy L, Barros CK, Procianoy RS. Incidence and risk factors for retinopathy of prematurity in very low and in extremely low birth weight infants in a unit-based approach in southern Brazil. *Eye (Lond)* 2009;23:25–30.
9. Gilbert C, Fielder A, Gordillo L, Quinn G, Semiglia R, Visintin P, et al. Characteristics of infants with severe retinopathy of prematurity in countries with low, moderate, and high levels of development: Implications for screening programs. 2005;115:e518–525.
10. Section on Ophthalmology American Academy of Pediatrics; American Academy of Ophthalmology; American Association for Pediatric Ophthalmology and Strabismus. Screening examination of premature infants for retinopathy of prematurity. 2006;117:572–576.
11. Early Treatment for Retinopathy of Prematurity Cooperative Group. Revised indications for the treatment of retinopathy of prematurity: Results of the early treatment for retinopathy of prematurity randomized trial. *Arch Ophthalmol.* 2003;121:1684–1694.
12. Ho SF, Mathew MR, Wykes W, Lavy T, Marshall T. Retinopathy of prematurity: An

optimum screening strategy. J AAPOS. 2005;9:584–588.

13. Larsson E, Holmström G. Screening for retinopathy of prematurity: Evaluation and modification of guidelines. Br J Ophthalmol. 2002;86:1399–1402.

14. Maheshwari R, Kumar H, Paul VK, Singh M, Deorari AK, Tiwari HK. Incidence and risk factors of retinopathy of prematurity in a tertiary care newborn unit in New Delhi. Natl Med J India. 1996;9:211–214.

15. Gupta VP, Dhaliwal U, Sharma R, Gupta P, Rohatgi J. Retinopathy of prematurity –Risk factors. Indian J Pediatr. 2004;71:887–892.

16. Palmer EA, Flynn JT, Hardy RJ, Phelps DL, Phillips CL, Schaffer DB, *et al.* Incidence and early course of retinopathy of prematurity. Ophthalmology 1991; 98: 1628-1640.

17. Vinekar A, Dogra M, Sangtam T, Narang A, Gupta A. Retinopathy of prematurity in Asian Indian babies weighing greater than 1250 grams at birth: ten year data from a tertiary care center in a developing country. Indian J Ophthalmol 2007; 55:331-336.

18. Jalali S, Anand R, Kumar H, Dogra MR, Azad RV, Gopal L. Programme planning and screening strategy in retinopathy of prematurity. Indian J Ophthalmol 2003; 51: 89-99.

19. Ng EY, Connolly BP, McNamara JA, Regillo CD, Vander JF, Tasman W. A comparison of laser photocoagulation with cryotherapy for threshold retinopathy of prematurity at 10 years. Part 1-Visual function and structural outcome. Ophthalmology 2002; 109; 928-934.

20. Connolly BP, Ng EY, McNamara JA, Regillo CD, Vander JF, Tasman W. A comparison of laser photocoagulation with cryotherapy for threshold retinopathy at 10 years. Part 2-Refractive outcome. Ophthalmology 2002; 109: 936-941.